

RESEARCH ARTICLE

Teodora Staykova

Inter- and intra-population genetic variability of introduced silkworm (*Bombyx mori* L.) strains raised in Bulgaria**Authors' address:**

Department of Developmental Biology,
Faculty of Biology, Plovdiv University,
Plovdiv, Bulgaria.

Correspondence:

Toedora Staykova
Faculty of Biology
Plovdiv University
24, Tsar Assen Str.
4000 Plovdiv, Bulgaria
Tel.: +359 32 261549
e-mail: tstaykova@yahoo.com

Article info:

Received: 20 November 2012

Accepted: 5 April 2013

ABSTRACT

The genetic variability of four populations belonging to two introduced silkworm strains (*Bombyx mori* L.) of various origins has been studied using isoenzymic analysis of six enzyme systems. Nonspecific esterases, phosphoglucomutase, malate dehydrogenase, acid phosphatase, alkaline phosphatase and hexokinase from different tissue of larvae 5th instar have been analysed using PAGE. Polymorphism in six from a total of nine loci has been found. Inter- and intra-population differences have been ascertained expressed in different allele composition of the gene pool and different frequencies of alleles. A higher degree of inter-population variability has been reported on the acid phosphatase and a lower one – on the phosphoglucomutase.

Key words: *Bombyx mori* L., genetic variability, isoenzymes.

Introduction

The choice of parental strains in the selection of mulberry silkworm (*Bombyx mori* L.) was mainly based on the study of their biological and technological properties and their genealogical data. The assessment of the parental strains, however, should be complex and should be carried out simultaneously by different indexes. Finding suitable markers to study the genetic heterogeneity is important for the selection of this type with regard to increasing the adaptive potential and their productivity (Mirhoseini & Gholami, 2002). Using isoenzymes as markers makes it possible to study the genetic diversity of the mulberry silkworm as well as to differentiate the strains (Eguchi, 1995; Goldsmith, 1995; Etebari et al., 2005). The established genetic variability could be used also to study the gene flow and origin of the separate strains (Gamo & Ohtsuka, 1980).

The purpose of this study was to determine the degree of genetic variability of various populations of the two introduced strains of mulberry silkworm raised in Bulgaria using isoenzyme markers.

Materials and Methods

Four populations from two strains of mulberry silkworm (*Bombyx mori* L.) with different origin were studied:

Line 22 – uni-bivoltine strain introduced from Uzbekistan. The egg serosa color is gray, chorion color is white and eggs are sticky. The larvae are white in color with markings. The cocoons are white in color and oval-elongated in shape.

Mziuri 1 – uni-bivoltine strain introduced from Georgia. The egg serosa color is gray, chorion color is white and eggs are sticky. The larvae are white in color with high marking. The cocoons are white, elongated with low constriction (Petkov et al., 2006).

All individuals were nourished at a standard regime of silkworm breeding in Sericultural Experiment Station Vratza and Agricultural University - Plovdiv. According to locations of rearing, populations were marked as Line 22 Vr and Mziuri 1 Vr (from Vratza) and Line 22 Pv and Mziuri 1 Pv (from Plovdiv). On the fifth day of the fifth instar 50 - 81 larvae were selected randomly from each population.

The spectrum of nonspecific esterases (EST, EC 3.1.1), malate dehydrogenase (MDH, EC 1.1.1.37) and acid phosphatase (ACP, EC 3.1.3.2) from hemolymph and the spectra of phosphoglucomutase (PGM, EC 5.4.2.2) and hexokinase (HK, EC 2.7.1.1) from silk glands as well as the spectrum of alkaline phosphatase (ALP, EC 3.1.3.1) from gut of larvae were studied by means of 7.5% PAGE (Daevis, 1964). Isolation of the tissues and preparation of the samples was made according to Staykova et al. (2003) and Staykova

RESEARCH ARTICLE

et al. (2004). Method of Shaw & Prasad (1970) was used to visualize the nonspecific esterases and malate dehydrogenase. Methods of Spencer et al. (1964) and Eaton et al. (1966) were used to visualize the phosphoglucomutase and hexokinase, respectively. Acid phosphatase isoenzymes were visualized according to Staykova et al. (2010) and alkaline phosphatase according to Boyer (1961).

Allele frequencies, mean number of alleles per locus, proportion of polymorphic loci at the 99% level, observed (H_o) and expected (H_e) heterozygosity, deviation from the Hardy-Weinberg equilibrium and Wright's fixation index (F_{ST}) (Wright, 1965), were calculated using BIOSYS-1 (Swofford & Selander, 1981).

Results and Discussion

Four of the studied enzyme systems – nonspecific esterases, phosphoglucomutase, malate dehydrogenase and acid phosphatase showed polymorphism (Table 1). Alkaline phosphatase and hexokinase were monomorphic across all individuals of the studied populations.

The nonspecific esterases in the mulberry silkworm (*Bombyx mori* L.) hemolymph are coded by different genes (Bes A, B, D and E), showing polymorphism (Egorova et al., 1985; He, 1995; Staykova et al., 2003; Staykova, 2008). In the gene pool of all populations included in this study, we found a three-allele polymorphism in Bes B locus, in the presence of Bes B₁, B₂ and B₃ alleles (Table 1). Bes D locus was also polymorphic and presented with three alleles (Bes D₁, D₂ and D₃) in the gene pool of Line 22 Vr, Line 22 Pv and Mziuri 1 Pv. Only in the gene pool of Mziuri 1 Vr we established a fourth allele (Bes D₀). We observed also a three-allele polymorphism in the Line 22 Pv population by Bes E locus (Bes E₁, E₂ and E₀). In the gene pool of Line 22 Vr and Mziuri 1 Pv, this gene was presented by two alleles (Bes E₁ and E₀). In all populations, where we found polymorphism in Bes E locus, Bes E₀ allele showed the highest frequency, which was fixed in the gene pool of Mziuri 1 Vr. In all studied populations of the two strains, Bes A locus was monomorphic.

Staykova (2006, 2008) describes polymorphism at the phosphoglucomutase locus (Pgm A) with the strains of mulberry silkworm raised in Bulgaria. In the gene pool of the studied populations of Line 22 and Mziuri 1, we ascertained the presence of three alleles (Pgm A₁, A₂ and A₃). In three out of four populations, the highest frequency was the one of Pgm A₂ (Line 22 Pv, Mziuri 1 Vr and Mziuri 1 Pv).

We found three-allele polymorphism by Mdh A locus in Line 22 Pv (Mdh A₁, A₂ and A₃). In the gene pool of the rest populations, Mdh A₂ allele was fixed. Marcato et al. (1990) also describe monomorphism by this locus. Egorova & Nasirillaev (1993) reported the presence of polymorphism of malate dehydrogenase in hemolymph, which coincides with our findings in Line 22 Pv.

Polymorphism in codominant alleles at the locus, coding the acid phosphatase of *Bombyx mori* L. hemolymph (blood phosphatase - Bph) was reported for the first time by Yoshitake & Akiyama (1964). Later on, it was also described by some other authors (Eguchi et al., 1988; Staykova et al., 2010). The Bph locus was mapped in 23rd chromosome of the mulberry silkworm (Fujimori et al., 1984). In the present study we also ascertained polymorphism in Bph gene, which was presented with three alleles in the gene pool of Mziuri 1 Vr (Bph A, B and 0) and with two alleles in the gene pool of Line 22 Vr (Bph B and 0) and of Mziuri 1 Pv (Bph A and B). Bph B allele was fixed in the gene pool of Line 22 Pv (Table 1).

In the gene pool of all studied populations, we established monomorphism in the loci controlling the alkaline phosphatase (Alp A) and hexokinase (Hk A). Staykova & Ivanova (2012) also reported lack of inter-breed polymorphism in hexokinase locus.

In the present study we ascertained differences in the composition of the gene pools of the pairs of populations of the same strain. These differences were related to the different frequencies of ranging of separate alleles in polymorphic loci, as well as to the lack or fixation of some alleles – only with one of the populations in the pair (Table 1). For example, Bes E₂ allele rarely found in the gene pool of Line 22 Pv (frequency 0.067), was eliminated by the gene pool of Line 22 Vr. Mdh A₂ allele, frequently found in the gene pool of Line 22 Pv (frequency 0.683), was fixed in the gene pool of Line 22 Vr. Bes D₀ allele, which was found in Mziuri 1 Vr with no high frequency (0.218), was eliminated by the gene pool of Mziuri 1 Pv. Bes E₀ allele, which was frequently found in Mziuri 1 Pv (frequency 0.703), was fixed in the gene pool of Mziuri 1 Vr. The elimination of rare alleles and the fixation of frequently found alleles in the gene pools of the populations of the same strain is probably a result of the effect of genetic drift. The fixation of Bph B allele in the gene pool of Line 22 Pv and the lack of Bph 0 allele of the gene pool of Mziuri 1 Pv, could also be a result of genetic drift, related with the so called “founder effect”.

RESEARCH ARTICLE

Table 1. Allele frequencies in populations tested.

Locus	Populations			
	Line 22 Vr	Line 22 Pv	Mziuri 1 Vr	Mziuri 1 Pv
Bes A				
A ₁	1.0	1.0	1.0	1.0
Bes B				
B ₁	0.250	0.683	0.462	0.516
B ₂	0.208	0.117	0.500	0.281
B ₃	0.542	0.200	0.038	0.203
Bes D				
D ₁	0.181	0.533	0.141	0.516
D ₂	0.681	0.350	0.141	0.438
D ₃	0.139	0.117	0.500	0.047
D ₀	0	0	0.218	0
Bes E				
E ₁	0.181	0.400	0	0.297
E ₂	0	0.067	0	0
E ₀	0.819	0.533	1.0	0.703
PgmA				
A ₁	0.167	0.267	0.013	0.063
A ₂	0.389	0.467	0.577	0.547
A ₃	0.444	0.267	0.410	0.391
MdhA				
A ₁	0	0.200	0	0
A ₂	1.0	0.683	1.0	1.0
A ₃	0	0.117	0	0
Bph				
A	0	0	0.205	0.625
B	0.389	1.0	0.205	0.375
C	0	0	0	0
D	0	0	0	0
0	0.611	0	0.590	0
Alp A				
A ₁	1.0	1.0	1.0	1.0
Hk A				
A ₁	0	0	0	0
A ₂	1.0	1.0	1.0	1.0

The mean number of alleles per locus varied from 1.9 (for Line 22 Vr and Mziuri 1 Pv) to 2.1 (for Line 22 Pv) (Table 2). The estimated percentage of polymorphic loci was lowest in Mziuri 1 Vr (44.4%). The observed heterozygosity (H_o) varied from 0.119 (for Line 22 Pv) to 0.164 (for Line 22 Vr). With all analyzed populations the expected heterozygosity (H_e) by polymorphic loci was higher than the observed one.

There were significant deviations of genotype frequencies from Hardy-Weinberg expectations at most of the loci in most populations ($P < 0.05$). Chi-Square (df: 1-6) tests showed that the deviations were generally in favour of homozygotes. The lower degree of observed heterozygosity and the higher degree of homozygotes proved the inbreeding effect.

RESEARCH ARTICLE

Table 2. Mean number of alleles per locus, proportion of polymorphic loci, observed (H_o) and expected heterozygosity (H_e)

Populations	Mean sample size per locus	Mean No. of alleles per locus	Percent polymorphic loci (P=0.99)	H_o	H_e
Line 22 Vr	50.0±0.0	1.9±0.30	55.6	0.164±0.059	0.279±0.094
Line 22 Pv	81.0±0.0	2.1 ±0.40	55.6	0.119±0.045	0.308±0.099
Mziuri 1 Vr	50.0±0.0	2.0±0.40	44.4	0.157±0.067	0.255±0.102
Mziuri 1 Pv	80.0±0.0	1.9±0.30	55.6	0.142±0.059	0.292±0.094

The estimated mean F_{ST} value was 0.1752 which shows that 17.52% of the overall genetic diversity observed was among populations. The highest value of intra-population genetic variability was calculated by Bph locus (0.3734) and the lowest one was calculated by Pgm locus (0.0324).

The results obtained in this study showed that in the studied populations of the two introduced strains Line 22 and Mziuri 1, there is an inter- and intra-population polymorphism at six enzyme loci (Bes B, Bes D, Bes E, Pgm, Mdh, Bph) and monomorphism at three loci (Bes A, Alp and Hk). The acid phosphatases from the hemolymph of the mulberry silkworm are very suitable markers to analyze the intra-population polymorphism. Apart from this group of isoenzymes, malate dehydrogenase and nonspecific esterases could also be used for the same purpose. Phosphoglucumutase is more suitable to study the inter-population polymorphism and determination of the degree of population genetic variability. The genetic variability determined in the course of this study is important for the analysis of separate populations and strains in order to use them in selection programs.

References

- Boyer SH. 1961. Alkaline phosphatase in human sera and placenta. Science, 134: 1002-1004.
- Daavis B. 1964. Disc electrophoresis. Methods and application to human serum proteins. Ann. N. Y. Acad. Sci., 121(4): 404-427.
- Eaton GM, Brewe GJ, Tashian RE. 1966. Hexokinase isozyme patterns of human erythrocytes and leucocytes. Nature, 212(5065): 944-946.
- Egorova T, Naletova E, Nasirillaev Y. 1985. Polymorphic system of silkworm haemolymph esterases as a criterion to make programs for parental specimens crossing. Biochemistry of Insects - Moscow, 54-62 (in Russian).
- Egorova T, Nasirillaev Y. 1993. Polymorphic enzymes of silkworm and their using in breeding. State committee of science and technics of republic Uzbekistan, GFNTI: 120 p. (in Russian)
- Eguchi M, Takahama Y, Ikeda M, Horii S. 1988. A novel variant of acid phosphatase isozyme from hemolymph of the silkworm, *Bombyx mori*. Japan Journal of Genetics, 63(2): 149-157.
- Eguchi M. 1995. Alkaline phosphatase isozymes in insects and comparison with mammalian enzyme. Comparative Biochemistry and Physiology, 111(2): 151-162.
- Etebari K, Mirhoseini SZ, Matindoost L. 2005. A study on interspecific biodiversity of eight groups of silkworm (*Bombyx mori*) by biochemical markers. Insect Science, 12(2): 87-94.
- Fujimori H, Shimizu K, Enokijima M. 1984. Chromosome mapping of the blood acid phosphatase gene in the silkworm, *Bombyx mori*. Journal of Sericultural Science of Japan, 53(2): 181-182.
- Gamo T, Ohsuka Y. 1980. Phylogenetic studies on the racial differentiation of the silkworm, *Bombyx mori* on the basis of polymorphic genes in haemolymph proteins. Bull. Sericul. Exp. Sta., 28(1): 15-50.
- Goldsmith MR. 1995. Genetics of the silkworm: revisiting an ancient model system. In: Goldsmith MR & Wilkins AS (eds), Molecular Model Systems in the Lepidoptera, Cambridge University Press, New York, p. 21-76.
- He JL. 1995. Studies on the inheritance of a deletion type of the blood esterase isoenzymes in the silkworm, *Bombyx mori* L. Sericologia, 35(1): 17-24.
- Marcato S, Trevisan M, Cappelozza L, Bisol P. 1990. Variabilita di sistemi gene-enzima, in razze di *Bombyx mori* (Lepidoptera, Bombycidae) utilizzate per la produzione di seta. Estratto da Redia, LXXIII (2): 595-608. (in Italian, English summary)
- Mirhoseini SZ, Gholami MR. 2002. Study on some economic characters of single, three-way and double crosshybrids obtained from four Iranian silkworm varieties. XIXth Congress of The International Sericultural Commission Proceedings, Thailand, p.190-194.
- Petkov N, Tzenov P, Petkov Z, Natcheva Y, Vasileva Y. 2006. Silkworm, *Bombyx mori* L. germplasm resources in Bulgaria. National Centre for Agrarian Sciences – Sofia, Sericultural Experiment Station – Vratza, p.264.
- Shaw C, Prasad R. 1970. Starch gel electrophoresis of enzymes – a compilation of Recipes. Biochemistry Genetics, 4(2): 297-320.
- Spencer N, Hopkinson D, Harris H. 1964. Phosphoglucumutase polymorphism in man. Nature, 204: 742-745.

RESEARCH ARTICLE

- Staykova T, Grekov D, Panayotov M. 2004. Electrophoretic analysis of nonspecific esterases in silkworm (*Bombyx mori* L.) female genital organs and eggs. *International Journal of Industrial Entomology*, 9(1): 59-63.
- Staykova T. 2006. Genetic control of phosphoglucomutase in mulberry silkworm (*Bombyx mori* L.). International Jubilee Scientific Conference "Problems of maintenance and utilization of mulberry and silkworm genetic resources", September 25 – 29, 2006, Vratza, Bulgaria, 197-201.
- Staykova T. 2008. Genetically-determined polymorphism of nonspecific esterases and phosphoglucomutase in eight introduced breeds of the silkworm, *Bombyx mori*, raised in Bulgaria. 8 pp. *Journal of Insect Science* 8:18, available online: insectscience.org/8.18.
- Staykova T, Ivanova E, Tzenov P, Vasileva Y, Arkova-Pantaleeva D, Petkov Z. 2010. Acid phosphatase as a marker for differentiation of silkworm (*Bombyx mori*) strains. *Biotechnology & Biotechnological Equipment*, 24(2SE): 379-384.
- Staykova T, Ivanova E. 2012. Inter-strain and intra-strain genetically-determined polymorphism of malate dehydrogenase and hexokinase and phylogenetic differentiation of the silkworm *Bombyx mori* L. University of Plovdiv "Paisii Hilendarski", Scientific studies – Biology, Jubilee collection "Biological sciences for more bright prospect", 253-263.
- Stoykova T, Popov P, Dimitrov B. 2003. Electrophoretic analysis of non-specific haemolymph esterases during silkworm (*Bombyx mori* L.) ontogenesis. *Sericologia*, 43(2): 153-162.
- Swofford DL, Selander RB. 1981. BIOSYS-1: A computer program for the analysis of allelic variation in genetics Rel. 1.0 Department of Genetics and Development University of Illinois at Urbana-Champaign, Urbana, Illinois 60801, USA.
- Wright S. 1965. The interpretation of population structure by F-Statistics with special regard to systems of mating. *Evolution*, 19(3): 395-420.
- Yoshitake N, Akiyama M. 1964. Genetical studies on the acid phosphatase in the blood of the silkworm, *Bombyx mori* L. *Japan Journal of Genetics*, 39: 26-30.