

## RESEARCH ARTICLE

Mahmoud M. Nour El-Dein<sup>1</sup>  
Abdou Eldyme A. Shereif<sup>2</sup>  
Fathey A. Mansour<sup>2</sup>  
Mohamed I. Abou-Dobara<sup>1</sup>  
Andrew S. Ball<sup>3</sup>

## Optimization of xylanase and peroxidase production from *Streptomyces* sp. K37

### Authors' addresses:

<sup>1</sup> Department of Botany,  
Faculty of Sciences, Damietta,  
Mansoura University, Egypt

<sup>2</sup> Department of Botany,  
Faculty of Sciences, Mansoura,  
Mansoura University, Egypt.

<sup>3</sup> Department of Biological Sciences,  
Essex University, England.

### Correspondence:

Mahmoud M. Nour El-Dein  
Department of Botany, Faculty of  
Sciences, Damietta, Mansoura  
University, Egypt.  
Tel.: +201094744952  
e-mail: mnoureldin@yahoo.com

### Article info:

Received: 9 September 2013

Accepted: 5 December 2013

### ABSTRACT

The optimal conditions for the production of xylanase and peroxidase from *Streptomyces* sp. K37 were investigated. The production of xylanase and peroxidase increased during the growth phase of the cultures after 72 hours. This indicates that the productions of such enzymes are wholly growth associated in these organisms. The optimum pH for xylanase and peroxidase production from *Streptomyces* sp. K37 occurred at pH 8.0 and pH 7.0. The optimum temperature for xylanase and peroxidase production by *Streptomyces* sp. K37 occurred at 50°C and 40°C respectively. Oat spelt xylan was the best carbon source for the production of xylanase and peroxidase from *Streptomyces* sp. K37. The optimum concentration of ball milled straw for the production of xylanase from *Streptomyces* sp. K37 was 1.0%, while the concentration for the production of peroxidase was 0.20%. The optimal concentration of yeast extract and casamino acid for the production of xylanase and peroxidase from *Streptomyces* sp. K37 was found to be 0.2%.

**Key words:** Characterization, production, xylanase, peroxidase, *Streptomyces* sp. K37

## Introduction

Xylans are one of the most abundant biopolymers, after cellulose, synthesised in the biosphere. In the cell wall of land plants, xylan is the most common hemicellulosic polysaccharide, representing more than 30% of the dry weight (Joseleau *et al.*, 1992). Xylans are composed of 1,4-linked  $\beta$ -D-xylopyranosyl residues (Whistler & Richards, 1970). Xylans from land plants are heteropolysaccharides that constitute a large group of related hemicelluloses showing a great variability in their structures (Joseleau *et al.*, 1992). This type of xylan contains different substituent groups in the backbone chain and in the side chain (Biely, 1985; Puls & Poutanen, 1989). The common substituent found on the backbone of xylan is acetyl, arabinosyl, and glucuronosyl residues (Whistler & Richards, 1970).

Lignin is considered to be the second most abundant natural polymer (after cellulose), making up 15 to 30% of the woody cells walls of gymnosperms (softwood) and

angiosperms (hardwood) (Gold & Alic, 1993). It forms a matrix surrounding the cellulose.

Lignin is a phenylpropanoid polymer synthesised from the phenolic precursor's coniferyl, synapyl, and p-coumaryl alcohols (Freudenberg, 1968; Sarkanen & Ludwig, 1971). Free-radical condensation of these precursors, initiated by plant cell wall peroxidases, results in the formation of a heterogeneous, amorphous, optically inactive, random, and highly branched polymer with at least 12 different types of linkages such as aryl ether and carbon-carbon bonds connecting the aromatic nuclei (Gold & Alic, 1993). These linkages are not subject to enzymatic hydrolysis. The majority of bonds within a typical lignin structure consist of aryl glycerol  $\beta$ -aryl ether ( $\beta$ -O-4-) and diaryl propane ( $\beta$ -1) bonds (Crawford, 1981). This unique structure requiring depolymerisation by extracellular oxidative mechanisms account for the recalcitrance of lignin toward degradation by most microorganisms (Crawford, 1981; Eriksson *et al.*, 1990; Gold *et al.*, 1989; Kirk & Farrell, 1987).

## RESEARCH ARTICLE

Yee *et al.* (1996) pointed out that since lignin also contains molecules of different chirality, lignin-degrading enzymes must be non-stereo selective (Crawford, 1981). Due to the irregular structure of the lignin, few organisms are able to degrade it. Sakakibara (1983) showed a schematic structural formula for lignin. Lignin is resistant to microbial degradation, although there are known lignin-degrading fungi, actinomycetes, and bacteria (Crawford 1981; Crawford & Crawford, 1984). This matrix retards the microbial depolymerisation of cellulose. The degradation of lignin represents a significant step in the global carbon cycle (Freudenberg, 1968; Sarkanen & Ludwig, 1971; Crawford, 1981). Also, lignin is an obstacle to the efficient utilisation of cellulose in a wide range of industrial processes.

Due to the variable composition and structure of xylans that can be obtained from different sources, the co-operative action of microbial enzymes that are excreted from the microorganism to affect the complete hydrolysis of the polysaccharide will be required. During the past five years some significant developments have given fresh impetus to the study of xylan-degrading enzymes (Coughlan & Hazlewood, 1993). These are the availability of recombinant DNA techniques for isolating single genes and expressing them in heterologous hosts devoid of hemicellulase activities, the introduction of biochemical techniques more suited to the analysis of complex mixtures of enzymes, and the commercial production of relevant synthetic and purified native substrates. A further important factor has been the general realisation that these enzymes have potentially valuable uses in the pulp and paper, textile and food industries and in agriculture.

For the complete hydrolysis of the xylan complex, number of enzymes is required. These enzymes work synergistically and co-operatively to convert xylan to its simple sugar. This complex enzyme system is common among bacteria (Woodward 1984; Wong *et al.*, 1988; Coughlan & Hazlewood, 1993). The hydrolysis of xylan is carried out by a number of main chain and side chain cleaving enzymes. The main chain enzymes include endo- $\beta$ -1,4-xylanases ( $\beta$ -1,4-D-xylanxylanohydrolase; EC 3.2.1.8),  $\beta$ -1,4-xylosidase ( $\beta$ -1,4-D-xyloside xylohydrolase; EC 3.2.1.37) and possibly exoxylanases ( $\beta$ -1,4-D-xylan xylohydrolase) while the side chain cleaving activities include  $\alpha$ -L-arabinofuranosidases and  $\alpha$ -D-glucuronodases as well as the esterases liberating acetyl, coumaroyl and feruloyl substituent's (Coughlan & Hazlewood, 1993).

$\beta$ -1,4-endoxylanase ( $\beta$ -1,4-D-xylanxylanohydrolase; EC 3.2.1.8) cleaves the internal glycosidic linkages of the heteroxylan backbone, resulting in a decreased degree of polymerisation (DP) of the substrate. During the early course of hydrolysis of xylan, the main products formed are xylooligosaccharides. As hydrolysis proceeds, these oligosaccharides will be further hydrolysed to xylotriose, xylobiose, and xylose (Dekker & Richards 1976, Wong *et al.* 1988, Debeire-Gosselin *et al.* 1992 a, 1992b).  $\beta$ -1,4-linkages between unsubstituted xylose residues and residues carrying monomeric glucuronosyl side-chains which can be cleaved by a novel enzyme, glucuronoxylanase (Nishitani & Nevins, 1991).

$\beta$ -D-xylosidase ( $\beta$ -1,4-D-xyloside xylohydrolase; EC 3.2.1.37) are exoglycosidases that hydrolyse short xylooligosaccharides and xylobiose from the non-reducing end to liberate xylose (Wong *et al.*, 1988). B-xylosidase is an important enzyme when complete hydrolysis of xylan is required (Biely, 1993). True  $\beta$ -xylosidases are able to cleave artificial substrates like p-nitrophenyl  $\beta$ -D-xyloside (Coughlan & Hazlewood, 1993). An important role of  $\beta$ -xylosidase seems to be relieving the end product inhibition of endoxylanase (Sunna & Antranikian, 1997).

Ferulic and p-coumaric acid are linked to xylan by ester bonds (Bacon *et al.*, 1975). Ferulic acid esterase (EC 3.1.1.-) cleaves the ester linkages between arabinose side chains and ferulic acids in xylan. Similarly, p-coumaric esterase (EC 3.1.1.-) cleaves the ester linkage between arabinose and p-coumaric acid (Müller-Harvey *et al.*, 1986; Mackenzie *et al.*, 1987).

Microbial xylan-degrading enzyme systems differ markedly with respect to whether they exist as free entities or as components of complexes (Hazlewood & Gilbert, 1993). Some microbial hemicellulases are secreted freely and do not associate with either the cell envelope or with other secreted proteins (Hazlewood *et al.*, 1992).

Sharma and Kumar (2013) pointed out that microbial endo-1,4- $\beta$ -xylanases are the preferred catalysts for xylan hydrolysis due to their high specificity, mild reaction conditions, negligible substrate loss, and side product generations.

Sunna and Antranikian (1997) identified the presence of a structure analogous to the cellosome, the xylanosome; as was reported in *Butyrivibrio fibrisolvens* H17c (Lin & Thomson, 1991). The extracellular complex B (C $\beta$ ) from *B. fibrisolvens* exists as a multisubunit protein aggregate. The complex has a

## RESEARCH ARTICLE

molecular weight higher than 669 kDa, and is composed of 11 protein bands with xylanase activity and 3 bands showing endoglucanase activity. The presence of xylanosomes in the anaerobic bacterium *Thermoanaerobacterium saccharolyticum* B6A-RI was reported by Zeikus *et al.*, (1991). In this bacterium, xylanase exists as a high molecular-weight complex, and enzyme activity is mainly cell-associated when the organism is grown on insoluble xylan. In the anaerobic thermophile *Clostridium thermocellum*, at least one well-characterised xylanase is a component of the highly-ordered multiprotein aggregate, termed the cellosome, which in early culture mediates the attachment of cells to cellulosic substrates (Grépinet *et al.*, 1988; Morag *et al.*, 1990).

The thermophilic actinomycete *Thermomonospora fusca* possess a complex multicomponent xylan-degrading system that consists of endoxylanase,  $\beta$ -xylosidase,  $\alpha$ -arabinofuranosidase, and acetyl esterase activities (Bachmann & McCarthy, 1991). The cooperative action between these enzymes has been studied.

Wong *et al.* (1988) noted that  $\beta$ -xylanases can be classified into two families according to their molecular weight in combination with their isoelectric point (pI). They have stated that one of the two families consists of high molecular weight enzymes with low pI, while the other consists of low molecular weight enzymes with high pI. This classification of  $\beta$ -xylanases according to Wong *et al.* (1988) was found to correspond with the classification of glycanases on the basis of hydrophobic cluster analysis and similarities in primary structures (Gilkes *et al.*, 1991; Shareck *et al.*, 1991). The high molecular weight  $\beta$ -xylanases belong to the F-family of glycanases while the low molecular weight  $\beta$ -xylanases belong to the G-family. Shareck *et al.* (1991) purified and sequenced three  $\beta$ -xylanases from *Streptomyces lividans* when growing this organism on xylan.

Furthermore, there is another classification of xylanases depending on the mode of action of the enzyme, i.e. the nature of the end product of the enzyme. Also  $\beta$ -xylanases can be classified into two groups according to the  $\alpha$ -L-arabinofuranosidase activity of  $\beta$ -xylanases. These two groups include arabinose-liberating  $\beta$ -xylanases and arabinose non-liberating  $\beta$ -xylanases (Dekker & Richards 1976; Reilly, 1981). The existence of arabinose liberating xylanases supports the suggestion that there could be genes encoding enzymes with two catalytic domains. One of these is responsible for  $\beta$ -xylanase and the other for  $\alpha$ -L-arabinofuranosidase activity (Matte & Forsberg, 1992).

The production of xylanases is common in nature. It has been reported to be found in some soil inhabiting organisms including bacteria, actinomycetes, fungi, rumen and marine bacteria, protozoa, insects and seeds of some terrestrial plants. Xylanases have been reported to occur in some thermophilic bacteria, actinomycetes and fungi, *Streptomyces thermoviolaceus* OPC-520 (Mackenzie *et al.*, 1987); *Nocardia* spp., *Saccharomonospora viridis* (Ball & McCarthy 1988, 1989); *Streptomyces lividans* 66 (Yasui *et al.*, 1988; Hernandez-Coronado *et al.*, 1997); *Thermomonospora fusca* BD25 (Bachmann & McCarthy, 1989; Irwin *et al.*, 1994); *Streptomyces chartreusis* GS 901 (Tsujiho *et al.*, 1992); *Streptomyces chattanoogensis* CECT 3336 (Kluepfel *et al.*, 1990); *Streptomyces* strains (Grabski & Jeffries, 1991; Grabski *et al.*, 1993); *Thermomonospora fusca* (Berens *et al.*, 1996); *Streptomyces* sp. strain B-12-2 (Senior *et al.*, 1992; Biely *et al.* 1993); *Streptomyces* sp. strain S38 (Elegir *et al.* 1994); *Streptomyces lividans* (Abd EL-Nasser & Foda 1995); *Streptomyces* spp. (Lopez-Fernandez *et al.*, 1998); *Streptomyces cyaneus* SN32 (Ninawe *et al.*, 2008); *Thermomonospora curvata* (Tuncer, 1999); *Microtetraspora flexuosa* SIIIX (Matsuo *et al.*, 2000), *Streptomyces roseiscleroticus* (Georis *et al.*, 2000).

Peroxidases catalyse the oxidation of a wide range of substrates including organic and inorganic compounds. They are widely found in plants, animals and microorganisms. Peroxidases have been detected intracellularly in bacteria (van Pée & Lingens, 1984, 1985), plants (Chibbar & Van Huystee 1984; Welinder 1985), oviparous animals (Wever *et al.*, 1980) and mammals (Moldoveanu *et al.*, 1982). Therefore, it can be concluded that peroxidases play an important role in biological systems. Extracellular peroxidases have also been described (Rob *et al.*, 1996, 1997). van Pée *et al.* (1987) studied a non-heme bromoperoxidase from *Streptomyces aureofaciens*. Extracellular peroxidases have also been described in other actinomycetes (Ramachandra *et al.*, 1988; Winter *et al.*, 1991; Godden *et al.*, 1992). They are secreted in multiple forms by a wide range of actinomycetes (Ball *et al.*, 1990; Trigo & Ball, 1994). It is expected that extracellular peroxidases have improved stability over their intracellular counterparts, particularly those from thermophiles (Iqbal *et al.*, 1994), and thus have potential for application in diagnostic kits (Mercer *et al.*, 1996).

Intracellular peroxidases of *Streptomyces* have also been specifically studied in relation to their role in the biosynthesis of halogenated antibiotics (van Pée *et al.*, 1987).

**RESEARCH ARTICLE**

An important difference between lignin peroxidase (LiP) and manganese peroxidase (MnP) is in the nature of the reducing substrate (Kirk & Farrell 1987; Gold *et al.*, 1989; Schoemaker, 1990; Gold & Alic, 1993). LiP catalyses the oxidation of non-phenolic lignin model compounds such as veratryl alcohol to veratryl aldehyde. Therefore the unique feature of this enzyme is that it is able to oxidise aromatic compounds with redox potential beyond that of horseradish peroxidase (HRP), while the oxidation of lignin and other phenols by MnP is dependent on free manganese ion (Glenn & Gold, 1985; Paszczyński *et al.*, 1986; Gold *et al.*, 1989).

*Streptomyces viridosporus* T7A produces a variety of enzymes that play a role in lignin degradation (Yee *et al.*, 1996). These extracellular enzymes include various peroxidases, cellulases, esterases, and endoglucanases (Crawford *et al.*, 1983, Borgmeyer & Crawford, 1985; Ramachandra *et al.*, 1987; Adhi *et al.*, 1989). One of the isoforms of peroxidase (AlIP-P3) produced by *Streptomyces viridosporus* T7A catalyses the degradation of lignin model compounds by oxidative cleavage of C $\alpha$ -C $\beta$  and C $\alpha$ -carbonyl bonds and is the only isoform detectable with an enzyme assay using 2,4-dichlorophenol as the substrate (Ramachandra *et al.*, 1988). Several peroxidases have been isolated, sequenced and characterised (Banci, 1997).

One of the most important applications of the peroxidase is the degradation of the harmful compounds (like chlorophenol) that can cause many problems to plant, animals, and humans. In addition to that, peroxidases play an important part in the degradation of lignin and lignocellulosic materials. Tuncer (1999) suggested that the ligninolytic activity could be used in a biotechnological application including treatment of bleach plant effluent (Michel Jr *et al.*, 1991) and the bioremediation of aromatic pollutants such as olive mill waste, a crucial problem in some Mediterranean countries (Sayadi & Ellouz, 1995).

The lignin peroxidase (LiP) of *Streptomyces viridosporus* T7A is known to degrade lignin and various phenolic compounds (Yee *et al.*, 1996). Therefore, *Streptomyces* may be used to degrade or mineralise recalcitrant waste compounds such as azo dyes (Paszczyński *et al.* 1992) and pesticides (Gauger *et al.*, 1986, Paszczyński *et al.*, 1992).

Xylanases can be used in different biotechnological applications. One of the most recent important applications is the use of xylanases in prebleaching of Kraft pulps for the paper industry. The bleaching of pulps depends on the removal of lignin from pulps and this essentially requires the use of toxic chemicals including chlorine and hypochlorite.

The use of xylanases in pulp bleaching and the formation of toxic organic chlorines have attracted public attention in the last few years (Viikari *et al.*, 1991, 1994; Coughlan & Hazlewood, 1993). The principle for the use of xylanases is that xylan hydrolysis facilitates the release of lignin and this minimises the need for chlorine and other toxic chemicals. The use of xylanases in pulp bleaching results in reduction the kappa number and increase of the brightness of the pulp. There are important aims for the process (Viikari *et al.*, 1991, 1994; Yang *et al.*, 1992; Daneault *et al.*, 1994).

Most of these applications will not require the presence of the purified enzymes. This is an important economic factor as the costs associated with purification of enzymes are high. However, in the case of pulp pre-bleaching a cellulases free xylanase is required. The presence of cellulases with xylanase in the process would destroy the cellulose fibres and that led to destroying of the pulp. Therefore, cellulases-free preparations are required. This may involve enzyme purification or the use of cellulases free strains such as *Streptomyces sp.* CH-M-1035 (Flores *et al.*, 1997).

The aim of this work is to optimize the conditions for xylanase and peroxidase production from *Streptomyces sp.* K37.

## Materials and Methods

### *Growth of isolates in liquid cultures*

The organisms were grown in conical flasks containing Sorensen medium (Sorensen, 1957) supplied with oat spelt xylan (1.0% (w/v) and yeast extract (0.0-1.2% (w/v)). Oat spelt xylan in some cases was replaced by different carbon sources or wheat straw. The pH of the medium was adjusted to 7.5 using NaOH and HCl (1.0 M) prior to sterilization. Flasks were inoculated and incubated at 50°C with shaking at 150 rev min<sup>-1</sup> for 48-72 hours.

### *Preparation of ball milled straw*

Wheat straw was cut into small pieces and milled in a rotating ball-milled apparatus (The Pascal Engineering Co. Ltd., Sussex, England). A fine powder of straw was obtained.

### *Preparation of inoculum*

A suspension of spores and hyphal fragments from pure colonies of the organisms from Petri dishes were used to inoculate flasks containing Sorensen medium (Sorensen, 1957). The pre-culture was incubated at 50°C at 150 rev min<sup>-1</sup> for 24-48 hours. Production flasks were then inoculated with an aliquot (1 ml) of this culture.

**RESEARCH ARTICLE*****Preparation of the crude enzymes***

After the incubation period, cultures were harvested by centrifugation at 12,000g for 10 minutes at 4°C using a Sorval® RC 5B Plus (Du Pont Company, Sorval® Instruments, Delaware, USA). Culture supernatants were used in assaying for extracellular enzyme activities (xylanase, cellulase and peroxidase). The remaining supernatants were used for the determination of intracellular protein. This was stored at -20°C for several weeks

***Determination of intracellular protein***

Harvested pellets were washed three times with distilled water. Washed pellets were centrifuged, washed and dissolved in 20 ml of NaOH (1M), and boiled for 20 minutes. Boiled pellets were allowed to cool and then centrifuged. Dilution of the clarified solution was used to determine the intracellular protein concentration of the organisms using the Bio-Rad protein assay (Bradford, 1976). In this assay, 800 µL of protein solution was mixed with 200 µL of Bio-Rad protein reagent and incubated at room temperature. After 5 minutes, the optical density at 595 nm was recorded using a micro plate reader, (Dias micro-plate reader Dynatech Laboratories Inc. USA and BioLinx 2.10™ software). Protein concentrations were determined from a standard curve with bovine serum albumin (BSA) ranging from 0 to 25 µg ml<sup>-1</sup>.

***Xylanase and cellulase activities determined using the Method of Miller (1959)***

Xylanase and cellulase activities were assayed by the estimation of reducing sugar released from oat spelt xylan and carboxymethylcellulose respectively. The reaction mixture contained the following: 200 µL of 1.0% oat spelt xylan or carboxymethylcellulose dissolved in 0.1 M potassium phosphate buffer (pH 7.0) and an aliquot of diluted culture supernatant or purified enzymes preparation. The mixture was incubated at 50°C for 20 minutes. The determination of reducing sugar released during the incubation mixture was detected by the dinitrosalicylic acid method of Miller (1959). Determination of the concentration of the reducing sugar was performed spectrophotometrically at 550 nm using the dinitrosalicylic acid (DNS) for the determination of reducing sugar. 200 µL of dinitrosalicylic acid reagent was added to 200 µL of the clarified reaction mixture and standards. After mixing, the mixtures were boiled for 5 minutes. After cooling under tap water, the optical density of the coloured product at 550 nm was taken using a Microplate reader. D-xylose and D-glucose standard

solutions were used to construct standard curves. Calibration curves constructed using D-xylose and D-glucose standards in the range of 0-5 µM ml<sup>-1</sup> were used. One unit of enzyme activity was defined as the amount of enzyme that released 1 µM of reducing sugar (expressed as xylose equivalents) per millilitre per minute under assay conditions. Enzyme and substrate controls were included routinely.

***Peroxidase assay***

This method was adapted from the assay developed by Ramachandra *et al.*, (1987, 1988). Peroxidase was assayed using 2,4-dichlorophenol (2,4-DCP) as substrate. The final reaction (1 ml) mixture contained potassium phosphate buffer (200 µL, 0.1 M, pH 7.0); 16 mM 4-aminoantipyrine (200 µL); 25 mM 2,4-dichlorophenol (200 µL); culture supernatant (200 µL) and 50 mM hydrogen peroxide (200 µL). The reaction was initiated by the addition of hydrogen peroxide which was added last. Samples were then incubated at 50°C for 1.0 minute. The increase in absorbance as a result of the oxidation of 4-aminoantipyrine was measured at 510 nm. This assay procedure was scaled down to a total volume of 200 µL in a microplate reader. One unit of enzyme activity was defined as the amount of enzyme required for an increase an absorbance of 1.0 unit ml<sup>-1</sup>.

***Environmental factors affecting xylanase and peroxidase production by Streptomyces sp. K37******Effect of incubation periods***

*Streptomyces sp. K37* was grown in liquid oat spelt xylan medium (Sorensen, 1957) supplemented with 2 g L<sup>-1</sup> yeast-extract as a basal medium. The medium was supplied with 10.0 g L<sup>-1</sup> oat spelt xylan. This medium was adjusted to pH 7.5 using 1 N NaOH and 1 N HCl, distributed into 100 ml conical flask containing 50 ml. Following inoculation with 1.0 ml of spore suspension, flasks were incubated at 50°C at 150 rev min<sup>-1</sup>. After 24, 48, 72, 96 and 120 hours, cell free extracts were obtained by centrifuging the bacterial filtrates at 12,000 g for 10 minutes at 4 °C. The cell free extracts were used for estimation the xylanase and peroxidase activities.

***Effect of different initial pH***

For studying the effect of initial pH on xylanase and peroxidase production, *Streptomyces sp. K37* was grown at 150 rev. min<sup>-1</sup> in liquid Sorensen medium supplemented with 2 g L<sup>-1</sup> yeast extract and 10.0 g L<sup>-1</sup> oat spelt xylan.

## RESEARCH ARTICLE

The pH was adjusted at 4.0, 5.0, 6.0, 7.0, 8.0, 9.0 and 10.0 using 1N NaOH or 1N HCl. After 72 hours incubation, the final pH was recorded, xylanase and peroxidase activities were determined as before.

*Effect of different temperatures*

*Streptomyces* sp. K37 was grown in liquid Sorensen medium supplied with 2 g L<sup>-1</sup> yeast extract and 10.0 g L<sup>-1</sup> Oat spelt xylan. Incubation was carried out at 30, 40, 50 and 6°C respectively and at 150 rev min<sup>-1</sup> for 72 hours. The supernatant was used for determination of xylanase and peroxidase activities.

*Effect of different carbon sources*

In this experiment, two types of carbon sources were used:

Type 1 - Oat spelt xylan, starch, phenylalanine, glycerol, cellulose and carboxymethylcellulose were added separately (1.0 %) (w/v) to the liquid Sorensen medium;

Type 2- Cellobiose and xylose (1.0%) (w/v) were added following sterilization by filtration through millipore filter (0.45 µm).

The pH was adjusted to 7.5 using 1N NaOH or 1N HCl. Flasks were inoculated and incubated at 50°C for 72 hours at 150 rev min<sup>-1</sup>. After incubation of *Streptomyces* sp. K37, the supernatant was used for estimation of xylanase and peroxidase activities.

*Effect of different concentrations of ball-milled straw*


*Streptomyces* sp. K37 was cultivated using Sorensen medium fortified by different concentrations of rice straw (0.00, 0.25, 0.50, 0.75, 1.00 and 1.25% (w/v)). The pH was adjusted to 7.5 by 1N NaOH and 1N HCl. Flasks were inoculated, incubated at 50°C and at 150 rev min<sup>-1</sup>. After 72 hours, cell free extracts were obtained by harvesting the cultures by centrifugation at 12,000 g for 10 minutes at 4°C. Cultures filtrates were used for estimation of xylanase and peroxidase activities.

*Effect of different concentrations of yeast extract and casamino acid*

*Streptomyces* sp. K37 was grown in Sorensen medium supplemented with different concentrations (0.0, 0.2, 0.4, 0.6, 0.8, 1.0, and 1.2 g L<sup>-1</sup>) of yeast extract and casamino acid. Cultures were incubated at 50°C at 150 rev min<sup>-1</sup> for 72 hours and pH 7.5.

**Results*****Growth and production of xylanase and peroxidase of Streptomyces sp. K37.***

Growth and production of xylanase and peroxidase of *Streptomyces* sp. K37 in Sorensen medium containing oat spelt xylan (1.0%) is shown in Figure 1. The production of xylanase and peroxidase increased significantly during the growth of the organism, with the maximum production of both enzymes detected after 72 hours (4.337 U/mg intracellular protein ml<sup>-1</sup> and 0.537 U/mg intracellular protein ml<sup>-1</sup> for xylanase and peroxidase respectively). Intracellular protein, used as a measure of cell biomass (1.459 U/mg intracellular protein ml<sup>-1</sup>) increased gradually and reached its maximum level after 72 hours of incubation. After 72 hours, the amount of the intracellular protein decreased until it reached a minimum level after 120 hours (0.163 U/mg intracellular protein ml<sup>-1</sup>).




**Figure 1.** Growth (▲), production of xylanase (◆) and peroxidase (■) by *Streptomyces* sp. K37 grown in Sorensen medium containing 1.0% oat spelt xylan. Data are represented as means of three replicates with standard errors.

***Effect of pH on the production of xylanase and peroxidase***

Maximum xylanase production occurred in the pH range of 6.0 and 8.0. Below pH 6.0, the activity of xylanase production increased with increasing pH until it reached the maximum production at pH 8.0 (8.813 U/mg intracellular protein ml<sup>-1</sup>). Above pH 8.0, xylanase yield decrease and

## RESEARCH ARTICLE

retained only 4.15% of the activity at pH 9.0. Below pH 7.0, peroxidase production increased with increasing of the pH of the medium and maximum peroxidase production occurred at pH 7.0 (0.927 U/mg intracellular protein ml<sup>-1</sup>). Above this pH, the yield of the enzyme decreased to 40% of its maximal activity at pH 8.0 and was completely inhibited at pH 10.0 (Figure 2).



**Figure 2.** Effect of pH on growth ( $\blacktriangle$ ) and production of xylanase ( $\blacklozenge$ ) and peroxidase ( $\blacksquare$ ) by *Streptomyces sp. K37* grown in Sorensen medium containing 1.0% (w/v) oat spelt xylan. Data are represented as means of three replicates with standard errors.


#### Effect of temperature on the production of xylanase and peroxidase

The optimum temperatures for the production of xylanase and peroxidase activities were 50°C and 40°C respectively. Peroxidase activity was reduced by 67.87% at 50°C. At 60°C, no xylanase and peroxidase could be detected (Figure 3).

#### Effect of carbon sources on the production of xylanase and peroxidase

The best carbon source for the enzyme activity was found to be oat spelt xylan (10.265 U/mg intracellular protein ml<sup>-1</sup> and 1.52 U/mg intracellular protein ml<sup>-1</sup> for xylanase and peroxidase, respectively). With regard to xylanase, phenylalanine, xylose, starch and carboxymethylcellulose were found to be good sources for the production of xylanase, with activity varying from 8.974 U/mg intracellular protein ml<sup>-1</sup> to 5.668 U/mg intracellular protein ml<sup>-1</sup>, while the growth on cellulose, glycerol and cellobiose resulted lower

activity ranging from 3.347 U/mg intracellular protein ml<sup>-1</sup> to 2.804 U/mg intracellular protein ml<sup>-1</sup>. Conversely, carboxymethylcellulose, glycerol and starch were found to be good substrates for the production of peroxidase with activity varying from 1.184 U/mg intracellular protein ml<sup>-1</sup> to 1.063 U/mg intracellular protein ml<sup>-1</sup>, while the organism produced low peroxidase activity when it grew on phenylalanine, xylose, cellobiose and cellulose, with activity varying from 0.423 U/mg intracellular protein ml<sup>-1</sup> to 0.027 U/mg intracellular protein ml<sup>-1</sup> (Figure 4a and 4b).




**Figure 3.** Effect of temperature on the production of xylanase ( $\blacklozenge$ ), intracellular protein ( $\blacktriangle$ ) and peroxidase ( $\blacksquare$ ) by *Streptomyces sp. K37* grown in Sorensen medium containing 1.0% (w/v) oat spelt xylan. Data are represented as means of three replicates with standard errors.

#### Effect of ball milled straw on the production of xylanase and peroxidase

The effect of different concentrations of ball milled straw on the production of xylanase and peroxidase was investigated (Figure 5). Straw was found to be a good source for xylanase production and activity increased gradually with the optimum concentration for production occurring at a ball milled straw concentration of 1.0% (w/v) (6.598 U/mg intracellular protein ml<sup>-1</sup>). Above this concentration, xylanase activity declined until it reached a minimum value at a ball milled straw concentration of 1.5% (w/v). However, the maximum peroxidase activity (0.109 U/mg intracellular protein ml<sup>-1</sup>) occurred at a concentration of 0.25%, after this, the production decreased and reaches the minimum activity at 1.5% (0.019 U/mg intracellular protein ml<sup>-1</sup>).


## RESEARCH ARTICLE



**Figure 4.** Effect of carbon sources on the production of xylanase (A) and peroxidase (B) by *Streptomyces sp.* K37 grown in Sorensen medium. Data are represented as means of three replicates with standard errors.

#### Effect of yeast extract and casamino acid on the production of xylanase and peroxidase

The effect of different concentrations of yeast extract and casamino acid on the production of xylanase and peroxidase by *Streptomyces sp.* K37 was also investigated (Figure 5). The highest xylanase production ( $10.265 \text{ U/mg intracellular protein ml}^{-1}$  and  $11.798 \text{ U/mg intracellular protein ml}^{-1}$ ) and peroxidase production ( $0.892 \text{ U/mg intracellular protein ml}^{-1}$  and  $2.315 \text{ U/mg intracellular protein ml}^{-1}$ ) were recorded in media supplemented with 0.2% yeast extract and casamino acid respectively. Above this concentration, xylanase and peroxidase production decreased significantly and reached the minimum level at a concentration of 1.2% ( $3.111 \text{ U/mg intracellular protein ml}^{-1}$  and  $2.142 \text{ U/mg intracellular protein ml}^{-1}$  for xylanase and  $0.230 \text{ U/mg intracellular protein ml}^{-1}$  and  $0.218 \text{ U/mg intracellular protein ml}^{-1}$  for peroxidase on yeast extract and casamino acids, respectively).



**Figure 5.** Effect of ball milled straw on the production of xylanase (◆), peroxidase (■) and intracellular protein (▲) by *Streptomyces sp.* K37 grown in Sorensen medium supplied with 0.2% yeast extract. Data are represented as means of three replicates with standard errors.

## Discussion

### Effect of incubation period on xylanase and peroxidase production

*Streptomyces sp.* K 37 produced high levels of xylanase and peroxidase when grown in Sorensen medium containing xylan as the main carbon source. Xylanase production from *Streptomyces sp.* K37 ( $1.06 \text{ U/ml}^{-1}$ ) and peroxidase production from *Streptomyces sp.* K37 were higher than that reported for *Streptomyces cyaneus* ( $0.15 \text{ U/ml}^{-1}$ ) (Wang et al., 1993), *Streptomyces badius* 252 ( $1.0 \text{ U/ml}^{-1}$ ), *Amycolata autotrophica* DSM 43099 ( $0.4 \text{ U/ml}^{-1}$ ) (Ball & McCarthy, 1989) and *Thermomonospora fusca* BD25 ( $0.52 \text{ U/ml}^{-1}$ ) (Tuncer, 1999).

Xylanase and peroxidase production by *Streptomyces sp.* K37, increased during the growth of the cultures in Sorensen medium with the maximum production detected after 72 hours incubation. After this period, the activities of the enzymes decreased. The observed peaking and troughing of the production of extracellular enzymes can be attributed to: (1) The differences in the timing of induction of separate components of the xylanase system; (2) The products of action of one component inducing the synthesis of another; (3) Differential inhibition by products of substrate hydrolysis; (4) Differential inactivation by proteases, or variation in the pH during cultivation conditions (Tuohy & Coughlan, 1992; Wang et al., 1993).

**RESEARCH ARTICLE**

A decrease in cellular protein was observed after 72 hours of growth of *Streptomyces* sp. K37. This probably resulted from cellular lysis, an observation previously reported by Godden *et al.* (1989).

These results indicate that the production of extracellular enzymes by *Streptomyces* sp. K37 was growth associated and this is in agreement with other actinomycetes (Ball & McCarthy, 1989; Trigo & Ball, 1994; Tuncer, 1999).

The production of peroxidase during growth of *Streptomyces* sp. K37 was similar to *Streptomyces avermitilis* UAH30. Rob *et al.* (1997) reported that the maximum specific peroxidase activity was obtained after 72 hours. In contrast, peroxidase production from *Streptomyces* sp. K37 was different from that reported by Iqbal *et al.*, (1994) who showed that the maximum production of peroxidase from *Streptomyces thermoviolaceus* was obtained after 50 hours. Lodha *et al.*, (1991) found that lignin peroxidase from *Streptomyces viridosporus* T7A was a non-growth associated, secondary metabolite with maximum lignin peroxidase activity occurring after 36 hours.

Production of extracellular peroxidase is common among actinomycetes and the enzyme plays an important role in the oxidation of a number of aromatic compounds and in the degradation of lignin. Peroxidase of *Streptomyces viridosporus* catalysed C-C bond cleavage of phenolic and nonphenolic arylglycerol- $\beta$ -aryl ether lignin model compounds (Ramachandra *et al.*, 1988). The maximal production of peroxidase by *Streptomyces* sp. K37 (0.35 U ml<sup>-1</sup>) was similar to that reported for *Streptomyces viridosporus* T7A (0.24 U/ml<sup>-1</sup>) (Ramachandra *et al.*, 1987), but higher than that reported for *Streptomyces thermoviolaceus* (0.04 U/mg) (Iqbal *et al.* 1994) and *Streptomyces avermitilis* (0.12 U/mg) (Rob *et al.*, 1997).

**Effect of pH on xylanase and peroxidase production**

Xylanase production was maximal at pH 8.0 for *Streptomyces* sp. K37 and the optimal production of peroxidase by *Streptomyces* sp. K37 was found to be at pH 7.0. These results are in accordance with the work that has been reported for *Streptomyces griseoflavus* and *Streptomyces violarius* (Abd EL-Nasser & Foda, 1995), for *Thermomonospora fusca* (Tuncer, 1999), for *Streptomyces badius* (Adhi *et al.*, 1989) and for peroxidase from *Streptomyces avermitilis* UAH30 (pH 7.5) (Rob *et al.*, 1997) and from *Streptomyces viridosporus* T7A (Lodha *et al.*, 1991) (pH 6.0).

When xylanase activity was expressed as U/mg intracellular protein, the comparative ability of the organism to produce xylanase was not changed, indicating that the results did not merely reflect differences in growth yield. This is an important point often overlooked when xylanase activities are reported (Ball & McCarthy, 1989). The authors also showed that xylanase production of *Streptomyces* strain EC1 and *Amycolata autotrophica* was growth associated in these organisms.

**Effect of temperature**

The optimal temperature for xylanase production occurred at 50°C for *Streptomyces* sp. K37 and the optimal production of peroxidase by *Streptomyces* sp. K37 was found to occur at 40°C. This is in accordance with the results obtained by Tuncer, (1999) and McCarthy (1987). The results of peroxidase production is in agreement with that observed by Rob *et al.* (1997) and Lodha *et al.*, (1991) who reported that the maximum production of peroxidase from *Streptomyces avermitilis* UAH30 and *Streptomyces viridosporus* T7A occurred at 45°C.

**Effect of carbon sources**

The best carbon source for xylanase production by *Streptomyces* sp. K37 was found to be birchwood xylan and oat spelt xylan. Low levels of xylanases were detected when the organisms were grown on xylose, starch, arabinose, sucrose, glucose, cellobiose, glycerol, phenylalanine, carboxymethylcellulose, and cellulose.

Abd EL-Nasser & Foda, (1995) found that lactose gave high levels of xylanase in cultures of *Streptomyces griseoflavus* and *Streptomyces violarius* while glucose utilisation resulted in complete suppression of enzyme formation. The endoxylanase activity from *Thermoanaerobacterium saccharolyticum* B6A-RI was catabolite repressed by glucose (Shao *et al.*, 1995).

In this work, xylanases of *Streptomyces* sp. K37 were induced by growth of the organisms on a medium containing xylan. This phenomenon has been observed for other actinomycetes (McCarthy *et al.*, 1985, Mackenzie *et al.*, 1987, Holtz *et al.*, 1991, Tuncer, 1999). Induction by xylan may be attributed to the role of xylobiose which, like cellobiose, is known to be potent inducer of xylanolytic enzymes in many microorganisms (Rho *et al.*, 1982). Wang *et al.*, (1992) characterised three xylanases from *Streptomyces cyaneus* and explained the induction mechanism of secretion for these enzymes. They stated that polymeric xylan is

**RESEARCH ARTICLE**

unlikely to penetrate the cells and the immediate inducers played an important role in the induction process. The same authors pointed out that the presence of xylan degradation products could be fortuitous, due to the presence of other cells in the environment that are already xylanolytic. They described an alternative hypothesis that the cell itself was able to sense the presence of xylan via such inducers and this may be as a result of the activity of xylanase that is synthesised constitutively.

The best carbon source for the optimal peroxidase production by *Streptomyces* sp. K37 was found to be oat spelt xylan and the organism was able to produce peroxidase when grown on xylose, carboxymethylcellulose, starch, cellulose, glycerol and phenylalanine. Lodha *et al.* (1991) reported that the production of peroxidase from *Streptomyces viridosporus* T7A was maximal when yeast extract was used as a carbon source and the production was reduced by 83% when yeast extract was replaced by carboxymethylcellulose.

***Effect of ball milled straw on the production of xylanase and peroxidase***

Ball milled straws were found to be good inducers of xylanase production. *Streptomyces* sp. K37 produced xylanases when they were grown in a medium containing straw. The optimum production for xylanase by *Streptomyces* sp. K37 production was found to occur in the range of concentrations between 1.25% to 1.50% ball milled straw. A similar observation was recorded by Wang *et al.* (1993) who reported that *Streptomyces cyaneus* produced nearly twice as much xylanase activity when grown on ball milled straw than when grown on xylan. Godden *et al.* (1989) reported that *Streptomyces* sp. isolated from compost produced higher levels of xylanase activity when grown on ball milled straw than xylan. Zimmermann *et al.* (1988) reported that the highest xylanase activities were produced by *Streptomyces* strains and the thermophile *Saccharomonospora viridis* when grown on ball milled straw. Furthermore, ground wheat straw was found to be a very good substrate for the production of xylanase by *Actinmadura* SIIx, producing a 1.5-fold increase in that activity levels compared with those produced during growth on xylan (Holtz *et al.* 1991). Similar results were recorded by Abd EL-Nasser & Foda, (1995) who reported that several agricultural by-products, including wheat bran, wheat straw and corn cobs, yielded higher xylanase activities than those produced on pure xylan by *Streptomyces griseoflavus* and *Streptomyces violarus*.

The production of the enzymes described in this work, during the growth of *Streptomyces* on agricultural by-products containing medium can be explained by the partial solubility of the components of agricultural material in water under the effect of high temperature and pressure of autoclaving. Part of the substrate becomes soluble and such material contains low molecular weight components as well as lignin carbohydrate derivatives. These components support the initial growth of organisms and the production of the enzymes.

*Streptomyces* sp. K37 produced low levels of peroxidase when grown in a medium containing ball milled straw as the main carbon source. This can be explained by the fact that the utilisation of xylan for peroxidase production proceeds much faster than the utilisation of ball milled straw, resulting in greater peroxidase production, during growth on xylan.

*Streptomyces* sp. K37 produced low levels of xylanase and peroxidase when grown in high concentrations of xylan and ball milled straw. This can be attributed to end product inhibition of simple sugars such as xylose, glucose and other oligosaccharides (Tuncer, 1999).

The production of microbial xylanases from cellulosic agro-industrial residues is of particularly interest because these residues are abundant, cheap and non-toxic, in addition to being underutilized (Stutzenberger, 1994; Patel & Ray, 1994).

***Effect of nitrogen sources. Effect of yeast extract and casamino acid***

*Streptomyces* sp. K37 produced high levels of xylanase and peroxidase when the medium was supplemented with 0.2% yeast extract and casamino acid, but high concentrations led to a decrease of xylanase and peroxidase production. This is in agreement with Yee *et al.* (1996) and Korus *et al.* (1991) who reported the highest production of lignin peroxidase from *Streptomyces viridosporus* T7A occurred using 0.3% (w/v) yeast extract, whereas higher concentrations inhibited peroxidase production, perhaps as a consequence of high concentrations of toxic compounds which have an inhibitory effect on growth and peroxidase production.

*Streptomyces* sp. K37 produced different levels of xylanase and peroxidase with varying C:N ratios and this may be explained as a different types of induction of enzymes. This may suggest different mechanisms of production of extracellular enzymes of *Streptomyces* sp. K37 when grown with different C:N ratios. *Streptomyces* sp. K37

## RESEARCH ARTICLE

produced high levels of extracellular enzymes when the C:N ratio was optimal and as the result of changing this ratio the organism produced low levels of extracellular enzymes. These results are in accordance with reported by Fernandez *et al.* (1995) and Tuncer (1999), who obtained different xylan degrading enzyme activities from *Streptomyces chattanoogensis* UAH 23 and *Thermomonospora fusca* BD25 when grown in media containing different C:N ratios, (4:1 and 1.3:1).

Two general areas of exploitation can be envisaged as a result of lignocellulose degradation: (i) production of biomass that can be used as a source of cheap single-cell protein or for mediating a secondary process; and (ii) release of monomers from polymeric plant material that can be used by a second organism for the growth and production of commercial biochemical's (Edwards, 1993).

## References

- Abd El Nasser NH, Foda MS. 1995. Formation physiology of xylanases by the grey series of *Streptomyces*. *Microbiol. Res.*, 150: 315-321.
- Adhi TP, Korus RA, Crawford DL. 1989. Production of major extracellular enzymes during lignocellulose degradation by two *Streptomyces* in agitated submerged culture. *Appl. Environ. Microbiol.*, 55: 1165-1168.
- Bachmann SL, McCarthy AJ. 1989. Purification and characterisation of a thermostable  $\beta$ -xylosidase from *Thermomonospora fusca*. *J. Gen. Microbiol.*, 135: 293-299.
- Bachmann SL, McCarthy AJ. 1991. Purification and cooperative activity of enzymes constituting the xylan-degrading system of *Thermomonospora fusca*. *Appl. Environ. Microbiol.*, 57 (8): 2121-2130.
- Bacon JSD, Gordon AH, Morris EJ, Farmer VC. 1975. Acetyl groups in cell wall preparations from higher plants. *Biochem. J.*, 149: 485-487.
- Ball AS, Godden B, Helvenstein P, Penninck M J, McCarthy AJ. 1990. Lignocarbhydrate solubilisation from straw by actinomycetes. *Appl. Environ. Microbiol.*, 56: 3017-3022.
- Ball AS, McCarthy AJ. 1988. Saccharification of straw by actinomycete enzymes. *J. Gen. Microbiol.*, 134: 2139-2147.
- Ball AS, McCarthy AJ. 1989. Production and properties of xylanases from actinomycetes. *J. Appl. Bacteriol.*, 66: 439-444.
- Banci L. 1997. Structural properties of peroxidases. *J. Biotechnol.*, 53: 253-263.
- Berens S, Kaspari H, Klemme JH. 1996. Purification and characterization of two different xylanases from the thermophilic actinomycete *Microtetraspora flexuosa* SIIX. *Antonie van Leeuwenhoek Int. J. Gen. & Mol. Microbiol.*, 69: 235-241.
- Biely P, Kluepfel D, Morosoli R, Shareck F. 1993. Mode of action of three endo-B-1,4-xylanases of *Streptomyces lividans*. *Biochimica et Biophysica Acta*, 1162: 246-254.
- Biely P. 1985. Microbial xylanolytic systems. *Trends in Biotechnology*, 3(11): 286-290.
- Biely P. 1993. Biochemical aspects of the production of microbial hemicellulases. In: *Hemicellulose and Hemicellulases* (Eds. Coughlan, M.P. and Hazlewood, G.P.) pp. 29-51, Portland Press, London.
- Borgmeyer JR, Crawford DL. 1985. Production and characterization of polymeric lignin degradation intermediate from two different *Streptomyces* spp. *Appl. Environ. Microbiol.*, 49(2): 273-278.
- Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal. Biochem.*, 72: 248-254.
- Chibbar RN, Van Huystee RB. 1984. Characterisation of peroxidase in plant cells. *Plant Physiol.*, 75: 956-958.
- Coughlan MP, Hazlewood GP. 1993.  $\beta$ -1,4-D-xylan-degrading enzyme systems: biochemistry, molecular biology and applications. *Biotechnol. Appl. Biochem.*, 17: 259-289.
- Crawford DL, Pometto AL, Crawford RL. 1983. Lignin degradation by *Streptomyces viridosporus*: isolation and characterization of a new polymeric lignin degradation intermediate. *Appl. Environ. Microbiol.*, 45(3): 898-904.
- Crawford DL. 1981. Microbial bioconversion of lignin to useful chemicals using a lignin-degrading *Streptomyces*. *Biotechnol. Bioengin. Symposium Series*, 11: 275-291.
- Crawford RL, Crawford DL. 1984. Recent advances in studies of the mechanisms of microbial degradation of lignins. *Enzyme Microb. Technol.*, 6: 434-442.
- Daneault C, Leduc C, Valade JL. 1994. The use of xylanase in kraft pulp bleaching. *Tappi J.*, 77: 125-131.
- Debeire-Gosselin M, Loonis M, Samain E, Debeire P. 1992a. Purification and properties of a 22 kDa endoxylanase excreted by a new strain of thermophilic *Bacillus*. In: *Xylans and Xylanases*. (Eds. Visser, J., Beldman, G., Kusters-van Someren, M.A. And Voragen, A.G. J.), pp. 463-466. Elsevier, Amsterdam.
- Debeire-Gosselin M, Loonis M, Samain E, Debeire P. 1992b. Isoxylanases from the thermophilic *Clostridium thermolacticum*. In: *Xylans and Xylanases*. (Eds. Visser, J., Beldman, G., Kusters-van Someren, M.A. And Voragen, A.G.J.), pp. 471-474. Elsevier, Amsterdam.
- Dekker RFH, Richards GN. 1976. Hemicellulases: their occurrence, purification, properties, and mode of action. *Adv. Carbohydr. Chem. Biochem.*, 32: 277-352.
- Edwards C. 1993. Isolation properties and potential applications of thermophilic actinomycetes. *Appl. Biochem. Biotechnol.*, 24: 161-179.
- Elegir G, Szakacs G, Jeffries TW. 1994. Purification, characterization and substrate specificities of multiple xylanases from *Streptomyces* sp. strain B-12-2. *Appl. Environ. Microbiol.*, 60(7): 2609-2615.
- Eriksson K-EL, Blanchette RA, Ander P. 1990. Biodegradation of lignin, In: *Microbial and enzymatic degradation of wood and wood components*. pp. 225-333. Springer-Verlag KG, Berlin.
- Fernández LCL, Rodríguez J, Soliveri J, Copa-Patinà JL, Perez-Leblic MJ, Arias ME. 1995. The effect of culture media on the production of xylan-degrading enzymes by *Streptomyces chattanoogensis* UAH 23. *J. Basic Microbiol.*, 35: 405-412.
- Flores ME, Pérez R, Huitrón C. 1997. B-xylosidase and xylanase characterisation and production by *Streptomyces* sp. CH-M-1035. *Lett. Appl. Microbiol.*, 24: 410-416.

## RESEARCH ARTICLE

- Freudenberg K. 1968. The constitution and biosynthesis of lignin, pp. 47-122. In: Constitution and biosynthesis of lignin. (Eds. Neish, A.C. and Freudenberg, K.), Springer-Verlag, New York.
- Gauger WK, MacDonald JM, Adrian NR, Matthees DP, Walgenbach DD. 1986. Characterisation of Streptomycetes growing on organophosphate and carbamate insecticides. *Arch. Environ. Contam. Toxicol.*, 15: 137-141.
- Georis J, Giannotta F, DeBuyl E, Granier B, Frere JM. 2000. Purification and properties of three endo-beta-1,4-xylanases produced by *Streptomyces* sp. Strain S38 which differ in their ability to enhance the bleaching of kraft pulps. *Enzyme Microbiol. Technol.*, 26 (2-4): 178-186.
- Gilkes NR, Henrissat B, Kilburn DG, Miller RC Jr, Warren RAJ. 1991. Domains in microbial  $\beta$ -1,4-glycanases: sequence conservation, function and enzyme families *Microbiol. Rev.*, 55: 303-315.
- Glenn JK, Gold MH. 1985. Purification and characterisation of an extracellular Mn (II)-dependent peroxidase from the lignin-degrading basidiomycete, *Phanerochaete chrysosporium*. *Arch. Biochem. Biophys.*, 242: 329-341.
- Godden B, Ball AS, Helvenstein P, McCarthy AJ, Penninickx MJ. 1992. Towards elucidation of the lignin degradation pathway in actinomycetes. *J. Gen. Microbiol.*, 138: 2441-2448.
- Godden B, Legon T, Helvenstein P, Penninickx M. 1989. Regulation of the production of hemicellulolytic and cellulolytic enzymes by a *Streptomyces* sp. growing on lignocellulose. *J. Gen. Microbiol.*, 135: 285-292.
- Gold MH, Alic M. 1993. Molecular biology of the lignin-degrading basidiomycete *Phanerochaete chrysosporium*. *Microbiol. Rev.*, 57: 605-622.
- Gold MH, Wariishi H, Valli K. 1989. Extracellular peroxidases involved in lignin degradation by the white rot basidiomycete *Phanerochaete chrysosporium*. *ACS Symp. Ser.*, 389: 127-140.
- Grabski AC, Forrester TI, Patel R, Jeffries TW. 1993. Characterization and N-terminal amino acid sequences of B-1,4-Endoxylanases from *Streptomyces roseiscleroticus*: Purification Incorporating a Bioprocessing Agent. *Protein Expression and Purification*, 4: 120-129.
- Grabski AC, Jeffries TW. 1991. Production, purification and characterisation of  $\beta$ -(1,4)-endoxylanase of *Streptomyces roseiscleroticus*. *Appl. Environ. Microbiol.*, 57: 987-992.
- Grépinet O, Chebrou M-C, Béguin P. 1988. Nucleotide sequence and deletion analysis of the xylanase gene (*xyn Z*) of *Clostridium thermocellum*. *J. Bacteriol.*, 170: 4582-4588.
- Hazlewood GP, Gilbert HJ. 1993. Molecular biology of hemicellulases In: Hemicellulose and Hemicellulases (Eds. Coughlan, M. P. and Hazelwood, G. P.) pp.103-126. Portland Press, London.
- Hazlewood GP, Lurie JI, Ferreira LMA, Gilbert HJ. 1992. *Pseudomonas fluorescens* subsp. Cellulose: an alternative model for bacterial cellulase. *J. Appl. Bacteriol.*, 72 (3): 244-251.
- Hernandez-Coronado MJ, Hernandez M, Centenera F, Perez-Leblic MI, Ball AS, Aries ME. 1997. Chemical characterisation and spectroscopic analysis of the solubilisation products from wheat straw produced by *Streptomyces* strains grown in solid-state fermentation *Microbiol.*, 143: 1359-1367.
- Holtz C, Kaspari H, Klemme JH. 1991. Production and properties of xylanases from thermophilic actinomycetes. *Antonie van Leeuwenhoek*, 59: 1-7.
- Iqbal M, Mercer DK, Miller PGG, McCarthy AJ. 1994. Thermostable extracellular peroxidases from *Streptomyces thermoviolaceus*. *Microbiol.*, 140: 1457-1465.
- Irwin D, Jung ED, Wilson BD. 1994. Characterization and sequence of a *Thermomonospora fusca* xylanase. *Appl. Environ. Microbiol.*, 60 (3): 763-770.
- Joseleau JP, Comtat J, Ruel K. 1992. Chemical structure of xylans and their interaction in the plant cell walls. In: *Xylans and Xylanases*. (Eds. Visser, J., Beldman, G., Kusters-van Someren, M. A. and Voragen, A. G. J.), pp. 1-15. Elsevier, Amsterdam.
- Kirk TK, Farrell RL. 1987. Enzymatic "combustion": the microbial degradation of lignin. *Ann. Rev. Microbiol.*, 41: 465-505.
- Kluepfel D, Vats-Mehta S, Aumont F, Shareck F, Morosoli R. 1990. Purification and characterization of a new xylanase (xylanase B) produced by *Streptomyces lividans* 66. *Biochem. J.*, 267: 45-50.
- Korus RA, Lodha SJ, Adhi TP, Crawford DL. 1991. Kinetics of peroxidase production by *Streptomyces viridosporus* and recombinant *Streptomyces lividans*. *Biotechnol. Prog.*, 7: 510-515.
- Lin LL, Thomson JA. 1991. An analysis of the extracellular xylanases and cellulases of *Butyrivibrio fibrisolvens* H17c. *FEMS Microbiol. Lett.*, 84: 197-204.
- Lodha SJ, Korus RA, Crawford DL. 1991. Synthesis and properties of lignin peroxidase from *Streptomyces viridosporus* T7A. *Appl. Biochem. Biotechnol.*, 28/29: 411-420.
- López-Fernandez CL, Rodriguez J, Ball AS, Copa-Patino JL, Perez-Leblic MI, Arias ME. 1998. Application of the affinity binding of xylanases to oat-spelt xylan in the purification of endoxylanase CM-2 from *Streptomyces chattanoogensis* CECT 3336. *Appl. Microbiol. Biotechnol.*, 50 (2): 284-287.
- Mackenzie CR, Bilous D, Schneider H, Johnson KG. 1987. Induction of cellulolytic and xylanolytic enzyme systems in *Streptomyces* spp. *Appl. Environ. Microbiol.*, 53 (12): 2835-2839.
- Matsuo N, Kaneko S, Kuno A, Kobayashi H, Kusakabe I. 2000. Purification, characterisation and gene cloning of two alpha-L-arabinofuranosidases from *Streptomyces chartreusis* GS 901. *Biochem. J.*, 346 (pt1): 9-15.
- Matte A, Forsberg CW. 1992. Purification, characterization and mode of action of endoxylanases 1 and 2 from *Fibrobacter succinogenes* S85. *Appl. Environ. Microbiol.*, 58 (10): 157-168.
- McCarthy AJ, Peace E, Broda P. 1985. Studies on the extracellular xylanase activity of some thermophilic actinomycetes. *Appl. Microbiol. Biotechnol.*, 21: 238-244.
- McCarthy AJ. 1987. Lignocellulose degrading actinomycetes. *FEMS Microbiol. Rev.*, 46: 145-163.
- Mercer DK, Iqbal M, Miller PGG, McCarthy AJ. 1996. Screening actinomycetes for extracellular peroxidase activity. *Appl. Environ. Microbiol.*, 62(6): 2186-2190.
- Michel FC (Jr), Dass SB, Grulke EA, Reddy CA. 1991. Role of manganese peroxidase and lignin peroxidase of *Phanerochaete chrysosporium* in the decolorisation of Kraft bleach effluent. *Appl. Environ. Microbiol.*, 57: 2368-2375.
- Miller GL. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.*, 31: 426-428.
- Moldoveanu Z, Tenovuo J, Mestecky J, Pruitt KM. 1982. Human milk peroxidase is derived from milk leukocytes. *Biochimica et Biophysica Acta.*, 718: 103-108.

## RESEARCH ARTICLE

- Morag E, Bayer EA, Lamed R. 1990. Relationship of cellulosomal and noncellulosomal xylanases of *Clostridium thermocellum* to cellulose-degrading enzymes. *J. Bacteriol.*, 172 (10): 6098-6105.
- Müller-Harvey I, Hartley RD, Harris PJ, Curzon EH. 1986. Linkage of *p*-coumaryl and feruloyl groups to cell wall polysaccharides of barley straw. *Carbohydr. Res.*, 148: 71-85.
- Ninawe S, Kapoor M, Kuhad RC. 2008. Purification and characterization of extracellular xylanase from *Streptomyces cyaneus* SN32. *Bioresource Technol.*, 99: 1252-1258.
- Nishitani K, Nevins DJ. 1991. Glucuronoxylan xylanohydrolase: A unique xylanase with the requirement for appendant glucuronosyl units. *J. Biol. Chem.*, 266: 6539-6543.
- Paszczynski A, Huynh V-B, Crawford RL. 1986. Comparison of ligninase-I and peroxidase-M2 from the white-rot fungus *Phanerochaete chrysosporium*. *Arch. Biochem. Biophys.*, 244: 750-765.
- Paszczynski A, Pasti-Grigsby MB, Goszczynski S, Crawford RL, Crawford DL. 1992. Mineralisation of sulphonated azo dyes and sulphanic acid by *Phanerochaete chrysosporium* and *Streptomyces chromofuscus*. *Appl. Environ. Microbiol.*, 58: 3598-3604.
- Patel BN, Ray RM. 1994. Production and characterisation of xylanase from a *Streptomyces* species grown on agricultural wastes. *World J. Biotechnol.*, 10: 599.
- Puls J, Poutanen K. 1989. Mechanisms of enzyme hydrolysis of hemicelluloses (xylans) and procedures for determination of the enzyme activities involved. In: *Enzyme Systems For Lignocellulose Degradation*. (Ed. Coughlan, M.P.), pp 151-165. Elsevier, Applied Science.
- Ramachandra M, Crawford DL, Hertel G. 1988. Characterization of an extracellular lignin peroxidase of the lignocellulolytic actinomycete *Streptomyces viridosporus*. *Appl. Environ. Microbiol.*, 54 (12): 3057-3063.
- Ramachandra M, Crawford DL, Pometto AL. 1987. Extracellular enzyme activities during lignocellulose degradation by *Streptomyces* spp.: A comparative study of wild-type and genetically manipulated strains. *Appl. Environ. Microbiol.*, 53: 2754-2760.
- Reilly PJ. 1981. Xylanases: Structure and function. In: *Trends in Biology and Fermentations for Fuels and Chemicals* (Ed. Hollaender, A.), pp. 111-129, Plenum Press, New York.
- Rho D, Desrochers M, Jursak L, Griguez H, Defaye J. 1982. Induction of cellulase in *Schizophyllum commune*: *Thiobacillus* as a new inducer. *J. Bacteriol.*, 149: 47-53.
- Rob A, Ball AS, Tuncer M, Wilson MT. 1996. Thermostable novel non-haem extracellular glycosylated peroxidase from *Thermomonospora fusca* BD25. *Biotechnol. Appl. Biochem.*, 24: 161-170.
- Rob A, Hernandez M, Ball AS, Tuncer M, Arias ME, Wilson MT. 1997. Production and partial characterisation of extracellular peroxidase produced by *Streptomyces avermitili* UAH30. *Appl. Biochem. Biotechnol.*, 62: 159-174.
- Sakakibara A. 1983. Chemical structure of lignin related mainly to degradation products. In: *Recent Advances in Lignin Biodegradation Research*. (Eds. Higuchi, T., Chang, H.M. and Kirk, T.K.), pp.12-23. Uni. Publ. Tokyo.
- Sarkanen KV, Ludwig CH. 1971. Lignins. Occurrence, formation, structure and reactions. Wiley-Interscience, New York.
- Sayadi S, Ellouz R. 1995. Roles of lignin peroxidase and manganese peroxidase from *Phanerochaete chrysosporium* in the decolorization of olive mill waste waters. *Appl. Environ. Microbiol.*, 61: 1098-1103.
- Schoemaker HE. 1990. On the chemistry of lignin biodegradation. *Recl. Trav. Chim. Pays-Bas Neth.*, 109: 255-272.
- Senior DJ, Hamilton J, Bernierjr RL. 1992. Use of *Streptomyces lividans* xylanase for bleaching of kraft pulps. In: *Xylans and Xylanases*. (Eds. Visser, J., Beldman, G., Kusters-van-Someren, M.A. and Varagen, A.G.J.), pp. 555-558. Elsevier, Amsterdam, London, New York, Tokyo.
- Shao W, DeBlois S, Wiegel J. 1995. A high molecular weight, cell associated xylanase isolated from exponentially growing *Thermoanaerobacterium* sp. strain JW1SL-YS485. *Appl. Environ. Microbiol.*, 61 (3): 937-940.
- Sharek F, Roy C, Yaguchi M, Morosoli R, Kluepfel D. 1991. Sequences of three genes specifying xylanases in *Streptomyces lividans*. *Gene.*, 107: 75-82.
- Sharma M, Kumar A. 2013. Xylanases. *British Biotechnology J.* 3(1): 1-28.
- Sorensen H. 1957. Microbial decomposition of xylan. *Acta. Agric. Scand. Supplementum*, 1; pp. 86.
- Stutzenberger FJ. 1994. Extracellular enzyme production by *Thermomonospora curvata* grown on bagasse. *J. Ind. Microbiol. Biotechnol.*, 13: 35-42.
- Sunna A, Antranikian G. 1997. Xylanolytic enzymes from fungi and bacteria. *Critical Reviews in Biotechnology*, 17(1): 39-67.
- Trigo C, Ball AS. 1994. Production of extracellular enzymes during the solubilisation of straw by *Thermomonospora fusca* BD25. *Appl. Microbiol. Biotechnol.*, 41: 366-372.
- Tsujibo H, Miyamoto K, Kuda T, Minami K, Sakamoto T, Hasegawa T, Inamori Y. 1992. Purification, properties, and partial amino acid sequences of thermostable xylanases from *Streptomyces thermoviolaceus* OPC-520. *Appl. Environ. Microbiol.*, 58(1): 371-375.
- Tuncer M. 1999. A study of the enzymology of lignocellulose degradation by *Thermomonospora fusca* BD25. Ph.D. thesis, University of Essex, UK.
- Tuohy MG, Coughlan MP. 1992. Production of thermostable xylan degrading enzymes by *Talaromyces emersonii*. *Bioresource Technol.*, 39: 131-137.
- van Pée K-H, Lingens F. 1984. Detection of a bromoperoxidase in *Streptomyces phaeochromogens*. *FEBS Lett.*, 173: 5-8.
- van Pée K-H, Lingens F. 1985. Purification of bromoperoxidase from *Pseudomonas aureofaciens*. *J. Bacteriol.*, 161(3): 1171-1175.
- van Pée K-H, Sury G, Lingens F. 1987. Purification and properties of a nonheme bromoperoxidase from *Streptomyces aureofaciens*. *Biol. Chem. Hoppe-Seyler.*, 368: 1225-1232.
- Viihari L, Kantelinen A, Sundquist J, Linko M. 1994. Xylanases in bleaching: from an idea to the industry. *FEMS Microbiol. Rev.*, 13: 335-350.
- Viihari L, Sundquist J, Kettunen J. 1991. Xylanases enzyme promotes pulp bleaching. *Paper Timber*, 73: 384-389.
- Wang P, Ali S, Mason JC, Sims PFG, Broda P. 1992. Xylanases from *Streptomyces cyaneus*. In: *Xylans and Xylanases* (Eds. Visser, J., Beldman, G., Kusters-van Sorensen, M.A. and Voragen, A.G.J.), pp. 225-234. Elsevier, Amsterdam.
- Wang P, Mason JC, Broda P. 1993. Xylanases from *Streptomyces cyaneus*: their production, purification and characterization. *J. Gen. Microbiol.*, 139: 1987-1993.

## RESEARCH ARTICLE

- Welinder KG. 1985. Plant peroxidases, their primary, secondary and tertiary structures and their relation to cytochrome c peroxidase. *Eur. J. Biochem.*, 151: 497-504.
- Wever R, Hamers MN, Weening RS, Roos D. 1980. Characterisation of the peroxidase in human eosinophils. *Eur. J. Biochem.*, 108: 491-495.
- Whistler RL, Richards EL. 1970. Hemicelluloses. In: *The Carbohydrates*. (Eds., Pigman, W. and Horton, D.), vol. 2a. pp. 447-469. Academic Press, New York.
- Winter B, Fiechter A, Zimmermann W. 1991. Degradation of organochlorine in spent sulfate bleach plant effluents by actinomycetes. *Appl. Environ. Microbiol.*, 57: 2858-2863.
- Wong KKY, Tan LUL, Saddler JN. 1988. Multiplicity of  $\beta$ -1,4-xylanase in microorganisms: functions and applications. *Microbiol. Rev.*, 52: 305-317.
- Woodward J, 1984. Xylanases: functions, properties and applications. *Top. Enzyme Ferment. Biotechnol.*, 8: 9-30.
- Yang JL, Lu G, Eriksson K-EL. 1992. The impact of xylanase on bleaching of kraft pulps. *Tappi J.*, 75: 95-101.
- Yasui T, Marui M, Kusakabe I, Nakanishi K. 1988. Xylanases of *Streptomyces*. *Methods in Enzymology*, 160: 648-654.
- Yee DC, Jahng D, Wood TK. 1996. Enhanced expression and hydrogen peroxide dependence of lignin peroxidase from *Streptomyces viridosporus* T7A. *Biotechnol. Prog.*, 12: 40-46.
- Zeikus JG, Lee C, Lee Y-E, Saha BC. 1991. Thermostable saccharides: new sources, uses, and biodesigns. *ACS Symp. Ser.*, 460: 36-51.
- Zimmermann W, Winter B, & Broda P. 1988. Xylanolytic enzyme activities produced by mesophilic and thermophilic actinomycetes grown on graminaceous xylan and lignocellulose. *FEMS Microbiol. Lett.*, 55: 181-186.