

RESEARCH ARTICLE

Punit Kumar Khanna ¹
Ratna Chandra ¹
Arun Kumar ²
Nidhi Dogra ¹
Heena Gupta ¹
Gourav Gupta ¹
Vijeshwar Verma ¹

Development of microsatellite markers and their correlation with morphological and chemical markers in *Withania somnifera* (L.) Dunal

Authors' addresses:

¹ School of Biotechnology,
Shri Mata Vaishno Devi University,
Kakryal, Katra, Jammu and Kashmir,
182320 India.

² Biodiversity and Applied Botany,
Indian Institute of Integrative Medicine
(CSIR), Jammu, Jammu and Kashmir,
180001 India.

Correspondence:

Ratna Chandra
School of Biotechnology,
Shri Mata Vaishno Devi University,
Kakryal, Katra, Jammu and Kashmir,
182320 India.
Tel.: +91-01991-285691
e-mail: ratna.chandra@smvdu.ac.in

Article info:

Received: 12 March 2015

Accepted: 6 January 2016

ABSTRACT

The genetic variation and relationships among 14 *Withania* accessions were evaluated using morphological, chemical and Simple Sequence Repeat (SSR) markers. Wild accessions are more robust and better performing in morphological and chemical metabolite accumulation than cultivated one. The results revealed that out of fourteen, four primers showed distinct polymorphism, indicating the robust nature of microsatellites in revealing polymorphism. The banding pattern was recorded in the form of 0-1 data sheet which was analyzed using unweighted pair group method with arithmetic mean (UPGMA) based on Jaccard's similarity coefficient. The cluster analysis showed higher level of genetic variation among the accessions. Similarity coefficients ranged from 0.125 to 1. The dendrogram revealed 3 major distinct clusters. Higher range of similarity values for related genotypes using simple sequence repeats (SSR) provides greater confidence for the assessment of genetic diversity and relationships. The polymorphism information content (PIC) value for the SSR loci ranged from 0.0 to 0.40. Higher PIC values were associated with higher level of polymorphism. Results of this study showed a high degree of variation among analyzed accessions, indicating an important source of genetic diversity that can be used in future breeding programs.

Key words: *Withania somnifera*, morphological, chemical, markers, microsatellite

Introduction

Withania somnifera (L.) Dunal is an important medicinal plant belonging to the family Solanaceae and is commonly called Ashwagandha or Indian Ginseng. It is also known as Indian Winter Cherry. It is a valuable shrub in Ayurvedic medicine and is classified as a rasayana and acknowledged to increase longevity and vitality (Mishra *et al.*, 2000). It occurs in Sub-Himalayan tract up to Punjab, Nepal, Rajasthan, Sind, dried region of India, Eastern and Western ghats, Andaman and Nicobar Islands, Sri Lanka, Myanmar, Pakistan, Malaysia,

Thailand, Java, Asian and African tropics.

The roots of Ashwagandha are primarily used in Ayurvedic medicine, although both the leaves and seeds contain medicinal properties. The three main groups of active constituents are steroidal lactones, alkaloids, and flavonoids, in addition to several amino acids and iron. The presence of withanolides and sitoindosides occur rarely in nature and only to the family Solanaceae. Withaferin A and withanolide D are the two main withanolides contribute to the most of the biological actions of *Withania* (Matsuda *et al.*, 2001). The plant has been found useful in the treatment of burns, wounds, dermatological disorders, gastrointestinal

RESEARCH ARTICLE

diseases, dysfunctions of the respiratory system, asthma, bronchitis and cancer (Bone, 1996; Grierson & Afolayan, 1999).

It is a superior class herb with multiple benefits. The bruised leaves of this plant are used in the treatment of tumors and as an anti-inflammatory agent (Jayaprakasam *et al.*, 2003). The methanolic extracts of different parts of *W. somnifera* exhibit therapeutic potential against various types of cardiovascular problems and are also effective against hyperlipidemia, obesity, (Mary *et al.*, 2003) and aging. In the traditional system of medicine Ayurveda, this plant is claimed to have potent aphrodisiac rejuvenative and life prolonging properties. It has general animating and regenerative qualities and is used among others for the treatment of nervous exhaustion, learning ability, memory capacity, insomnia, tiredness potency issues and coughing.

Pharmacological activities of this plant includes physiologic and metabolic restoration, cognitive function improvement in geriatric states and recovery from neurodegenerative disorders like convulsions, tardive dyskinesia etc (Dhuley, 2000; Bhattacharya *et al.*, 2002). Other investigations indicated that *W. somnifera* also has antistress (Bhattacharya & Muruganandam, 2003) immunomodulatory, cytotoxic, antibacterial and antifungal (Atta-ur-Rahman *et al.*, 1998). This plant is reputed to promote vitality during recovery from chronic illnesses (Singh *et al.*, 2002) and is useful for pain management in arthritic conditions (Bandyopadhyay & Jha, 2003).

Analysis of genetic diversity in crop species is an important component of crop improvement programs and it is useful in constructing genetic maps. Study of genetic diversity is the process by which variation among individuals or groups of individuals or populations is analyzed by a specific method or a combination of methods (Mohammadi & Prasanna, 2003). Genetic diversity studies can be done at different levels such as morphological, cytological, biochemical and DNA marker systems. Molecular markers have been reported to be useful in genotype identification and introgression of desirable traits from diverse or wild germplasm into the available cultivars to broaden the genetic base (Thompson *et al.*, 1998). Molecular markers have proven useful for assessment of genetic variation in germplasm collections (Mohammadi & Prasanna, 2003).

SSRs are the markers of choice in many plant breeding programs because they are multi-allelic co-dominant markers, PCR-based, easily reproducible, randomly

and widely distributed along the genome and their analysis may be automated. Microsatellite polymorphism is based on the different numbers of a short repeated motif at a given locus (Rafalski *et al.*, 1996). Microsatellite sequences are mainly expressed as variation in the copy number of tandem repeats at a particular locus, especially suited to distinguish closely related genotypes; because of their high degree of variability, they are, therefore, favored in population studies (Smith & Devey, 1994) and for the identification of closely related cultivars (Vosman *et al.*, 1992). SSR markers are now considered to be among the most powerful genetic marker as they detect the variation in allele frequency at many unlinked loci. Simple sequence repeat (SSR) markers have been proposed as one of the most suitable markers for the assessment of genetic diversity of plants and animals. Expansion and contraction of SSR repeats in genes of known function can be tested for association with phenotypic variation or, more desirably, biological function (Ayers *et al.*, 1997). Several studies have found that genic SSRs are useful for estimating genetic relationship and at the same time provide opportunities to examine functional diversity in relation to adaptive variation (Eujay *et al.*, 2001; Russell *et al.*, 2004).

Development of SSR markers through genomic library construction involves enormous cost and time. However, identification of transferable markers from the related species has proved to be an important way to develop SSR markers and it was successfully achieved in the other crops (Peakall *et al.*, 1998; Somta *et al.*, 2009; Datta *et al.*, 2010 a,b; Datta *et al.*, 2011; Chandra, 2011). The aim of this study was to develop and characterize microsatellite markers in *Withania somnifera*.

Materials and Methods

Plant material and DNA isolation

Fourteen *Withania* accessions available at medicinal garden of Indian Institute of Integrative Medicine (IIIM), Jammu were used as experimental material (Table 1). Approximately 2 g of young leaf material of each was used to extract and purify genomic DNA through the protocol developed by Doyle & Doyle (1990) with slight modifications as described by Sharma *et al.* (2003). Approximate DNA yields were calculated by a UV-visible spectrophotometer and also quantified in 0.8% (w/v) agarose gels by comparison with standard 1 kb DNA ladder.

RESEARCH ARTICLE

Table 1. *Ashwagandha* germplasm selection and its source.

No.	Accession Number	Origin
1 ^a	AGB001	Manasa (M.P.)
2	AGB002	Bikaner (Rajasthan)
3	AGB003	Jammu (J&K)
4	AGB009	Amritsar (Punjab)
5	AGB012	Joginder Nagar (HP)
6 ^a	AGB015	Dabur Research Foundation (Ghaziabad)
7	AGB017	Lucknow (UP)
8	AGB019	Lucknow (UP)
9 ^a	AGB025	Neemuch (MP)
10	AGB030	Bhopal (MP)
11 ^a	AGB036	Pune (Maharashtra)
12	AGB042	Jodhpur (Rajasthan)
13 ^a	AGB053	Hyderabad (AP)
14 ^a	AGB055	Hyderabad (AP)

^a means cultivated accession, while other are wild.

Recording of morphological diversity

Based on 10 randomly selected plants from each plot, observations were recorded on following 13 traits: plant height (cm), plant dry biomass (gm), number of shoots plant⁻¹, leaf apex, leaf margin, leaf thickness, leaf hairyness, root type, root texture, root colour, root bark, dry root yield (gm), dry root yield hectare⁻¹ (quintal).

Chemical marker analysis

The examination of marker withanolides (withanolide A, withanone, withaferin A and 12-deoxy-withastramonolode) content in root was carried out by HPLC method (Khajuria *et al.*, 2004).

SSR analysis

All SSR primers were synthesized based on published sequence information (Benor *et al.*, 2008; Grover *et al.*, 2009). A total of 14 SSR primer sequences received from Integrated DNA Technologies, Faridabad, India. These primers were used for PCR amplification of *Withania* DNA. The polymerase chain reaction (PCR) was carried out in a final volume of 20 µl containing 10X reaction buffer 2 µl (50 mM KCl, 1.5 mM MgCl₂, 10 mM TrisHCl pH 9.0), dNTPs (1.25 µM each) 2 µl, forward microsatellite primer 0.15 µl (200 µM), reverse microsatellite primer 0.15 µl (200 µM), Taq polymerase (3 U µl⁻¹) 0.15 µl and

DNA 50-100 mg. PCR protocol consists of one cycle of a pre-denaturation step at 94°C followed by 35 amplification cycles (1 minute denaturation at 95°C, 1 minute annealing, 2 minute extension at 72°C) and ending with a final 10 minute elongation at 72°C. The PCR products were separated on 3% agarose gel in 1X TBE buffer containing 0.01% of ethidium bromide. Electrophoresis was carried out at a constant voltage of 80 V for 3 hrs and the amplified product was visualized under ultra-violet (UV) light and photographed using gel documentation system "Biorad". Annealing temperature was adjusted based on the specific requirement of each primer combinations. All the accessions were screened for presence/absence of SSR bands.

SSR data scoring and statistical analysis

All the accessions were scored for the presence (1) and absence (0) of the SSR bands and a binary matrix was obtained. These data were used to assess genetic relationships based on Jaccard's similarity index. Cluster analysis was conducted with the unweighted pair group method based on arithmetic averages to generate a dendrogram using free online software (<http://genomes.urv.cat/UPGMA>) by Garcia-Vallve *et al.* (1999).

Results

The fourteen *Withania* accessions were evaluated for plant height (cm), plant dry biomass (gm), number of shoots per plant, leaf apex, leaf margin, leaf thickness, leaf hairyness, root type, root texture, root colour, root bark, dry root yield (gm) and dry root yield per hectare (quintal). Variation in morphological characters of *Withania* accessions are summarized in Table 2. The wild accessions were very prolific based on quantitative traits of plant, leaf and root characters, viz., plant height (129–135 cm), plant dry biomass (65.4 gm), number of shoots per plant (6–7), root length (33–36.6 cm), root diameter (2.5–3.4 cm), dry root yield (15.6 gm) and dry root yield per hectare (10-12.4 quintal). Both wild and cultivated accessions portrayed a wide range of diversity in qualitative and quantitative characters however root texture, root colour and root bark were constant. Therefore morphologically, the wild accession could be separated easily from cultivated accessions.

The success of genetic improvement in this medicinal crop strongly depends on diversity of chemical metabolite

RESEARCH ARTICLE

content along with high potential for root yield. The accessions differed quantitatively for withanolides estimated by HPLC method. The total withanolide content ranged from 0.004 to 0.079% on dry root basis. The chemotypes with high content of specific compound are very important from pharmaceutical point of view. Chemical profiling of roots of wild population showed that withanolide A and withanone increases gradually with the advancement of age but decreases in tertiary roots afterwards. Both withanolide A and withanone in cultivated population have the same pattern of increase initially but decrease in contents in secondary roots, followed by increase in tertiary roots. Leaf is the site of synthesis of withaferin A and translocated later on to primary,

secondary and tertiary roots. Withaferin A is detected only in tertiary roots (0.004 gm) of wild type whereas in secondary (0.027 gm) and tertiary roots (0.018 gm) of cultivated type (Table 3). Interestingly, the wild accessions had the maximum withanolide-A (0.079 gm) in secondary roots and withanone (0.038 gm) in secondary roots.

DNA fingerprinting is a powerful tool as it provides sharply defined and repeatable genotypic descriptors than characterization based on morphological characteristics. SSR markers are a powerful tool for cultivar identification and analysis of genetic variation. In this study, fourteen microsatellite markers were used to test the genetic diversity of 14 accessions.

Table 2. Morphological characteristics of *Withania somnifera* (L.) Dunal.

Characteristics	Cultivated accession	Wild accession
Plant height (cm)	44-51 cm	129-135 cm
Plant dry biomass (gm)	32.4	65.4
No. of shoots per plant	4-5	6-7
Leaf apex	Obtuse tapering	Sharply acute
Leaf margin	Wavy	Entire
Leaf thickness	Subcoriaceous	Membranous
Leaf hairyness	Dense	Slight
Root type	Unbranched, erect, 11-16.8 cm long, 1.7 – 2.5 cm diameter	Unbranched, rarely branched at lower portion, 33-36.6 cm long, 2.5 – 3.4 cm diameter
Root texture	Very soft, tuberous	Soft, tuberous
Root colour	Creamish yellow	Light creamish
Root bark	Smooth, thin creamish colour	Smooth thick pale brown
Dry root yield (gm)	4.5	15.6
Estimated dry root yield Hectare ⁻¹ (quintal)	4-4.8	10-12.4

Table 3. Chemical profile of roots of wild and cultivated accessions based on 3 marker compounds ($g100g^{-1}$) at flowering stage.

Accession	Root type	Withanolide A	Withanone	Withaferin A
Cultivated	Primary	0.047	0.11	-
	Secondary	0.023	0.021	0.027
	Tertiary	0.030	0.031	0.018
Wild	Primary	0.054	0.19	-
	Secondary	0.079	0.038	-
	Tertiary	0.042	0.034	0.004

RESEARCH ARTICLE

Ten (71.42%) primers did not show polymorphism and were not considered for further analysis. The remaining four (28.57%) markers produced polymorphic banding pattern and it was possible to differentiate all the accessions used in the present study. The amplification band number differs from primer to primer, ranging from 2 to 9 (polymorphism band number: minimum of 2 and maximum of 9). Among all the selected primers, P27 and P31 have more amplification and polymorphism bands than the others. In order to demonstrate the potentiality of the SSR markers, PIC and H_e values were used as parameter, which varied with a range between 0-0.40. Our results indicate that markers with large number of alleles are informative for population studies. The high number of alleles per locus per plant in *Withania* could be due to heterozygosity and allogamous nature of cultivated *Withania*.




Figure 1. *Withania* genomic DNA isolated from different accessions: Lane (A) AGB-001, (B) AGB-002, (C) AGB-003, (D) AGB-009, (E) AGB-012, (F) AGB-015, (G) AGB-017, (H) AGB-019, (I) AGB-025, (J) AGB-030, (K) AGB-036, (L) AGB-042, (M) AGB-053 and (N) AGB-055.




Figure 2. Amplification profile of 14 accessions of *Withania* with P27 primer; M: Marker 1 kb ladder; Lanes 2-15: *Withania* accessions.

Basing on SSR markers an UPGMA analysis was performed. All 14 *Withania* accessions were discriminated successfully by SSR markers (Figure 3). The accessions were classified into three major clusters with Jaccard's similarity coefficient ranging from 0.125 to 1 (Table 5). The vast range of similarity coefficient between any two

accessions indicated the presence of wide genetic variability among the accessions studied. The first group gathered 3 cultivated accessions AGB015, AGB036 and AGB055, while Group II contains three accessions which involve AGB-001, AGB-025 and AGB-053. All the cultivated accessions were included in both the above groups. Group III was further divided into two subgroups (Subgroups 3a and 3b) containing 5 (AGB003 and AGB002; AGB019, AGB017 and AGB009) and 3 (AGB030, AGB042 and AGB012) accessions, respectively of wild group.




Figure 3. Dendrogram for the 14 accessions derived from cluster analysis of SSR data.

Simple sequence repeat (SSR) DNA markers have been shown to be highly polymorphic in soybean (Akkaya et al., 1992) used several types of SSRs to analyze the diversity of 43 soybean genotypes including ancestral and domestic cultivars representing the northern and southern U.S. gene pools. They determined that SSRs with (AT) and (ATT) repeat motifs were highly polymorphic in soybean and identified up to eight alleles at a single locus. Rongwen et al. (1995) identified 11 to 26 alleles at each of seven SSR loci in a diverse sample of soybean genotypes that included U.S. cultivars, *G. max* and *G. soja*. Maughan et al. (1995) detected 79 alleles across five SSR loci in a sample of 94 soybean accessions of *G. max* and *G. soja* genotypes. With 20 SSR markers, Diwan & Cregan (1997) were able to distinguish the 35 soybean genotypes that accounted for about 95% of the alleles present in North American soybean. They detected an average of 10.1 alleles per locus and an average marker diversity of 0.80.

SSR markers are ideal for distinguishing and identification between accessions that are genetically very similar (Saker et al., 2005). Their potential for automation and their inheritance in a codominant manner

RESEARCH ARTICLE

are additional advantages when compared to other types of molecular markers. Up to 37 different alleles for the one SSR locus have been found in barley (Saghai Maroof *et al.*, 1994). The Polymorphism Information Content (PIC) value depends on the diversity of the varieties tested (Pillen *et al.*, 2000; Hamza *et al.*, 2004).

Table 4. Simple sequence repeats (SSR) primers, their sequence and some information about generated bands in this study.

Primer name	Forward primer (5'-3') Reverse primer (5'-3')	Microsatellite motif	Allele size range (bp)	Scorable bands	PIC	H
P27	CCGCCTCTTCACTTGAAC CCAGCGATACGATTAGATACC	(CAG) ₇	130–175	9	0.370	0.365
P29	CGGGTGACACACGAGATTTT CCGCGAAAAGAAGTGGTAAG	(AG) ₇	150-162	2	0.0	0.0
P31	TGTGTTGCGTCATTACCACTAAAC CCCAACCACCAATACTTTCC	(AT) ₉	200-300	7	0.40	0.39
P32	TGGGTATGGGATTTACACCAA AAACGAAGGCAACAACGAAG	(AGA) ₂ (GAA) ₇	180-275	8	0.187	0.184

Table 5. Jaccard similarity matrix for 14 *Withania* accessions based on SSR data.

	AGB0 12	AGB0 42	AGB0 09	AGB0 55	AGB0 02	AGB0 53	AGB0 25	AGB0 36	AGB0 30	AGB0 03	AGB0 01	AGB0 17	AGB0 19	AGB0 15
AGB0 12	1	0.875	0.400	0.286	0.400	0.273	0.333	0.286	0.875	0.400	0.333	0.400	0.400	0.286
AGB0 42		1	0.500	0.250	0.500	0.250	0.417	0.250	1.000	0.500	0.417	0.500	0.500	0.250
AGB0 09			1	0.286	0.556	0.167	0.455	0.286	0.500	0.556	0.455	1.000	1.000	0.286
AGB0 55				1	0.125	0.125	0.222	1.000	0.250	0.125	0.222	0.286	0.286	1.000
AGB0 02					1	0.167	0.333	0.125	0.500	1.000	0.333	0.556	0.556	0.125
AGB0 53						1	0.600	0.125	0.250	0.167	0.600	0.167	0.167	0.125
AGB0 25							1	0.222	0.417	0.333	1.000	0.455	0.455	0.222
AGB0 36								1	0.250	0.125	0.222	0.286	0.286	1.000
AGB0 30									1	0.500	0.417	0.500	0.500	0.250
AGB0 03										1	0.333	0.556	0.556	0.125
AGB0 01											1	0.455	0.455	0.222
AGB0 17												1	1.000	0.286
AGB0 19													1	0.286
AGB0 15														1

Cophenetic correlation coefficient (CP): 0.95

RESEARCH ARTICLE

Discussion

On the basis of banding pattern, SSRs were effectively used for molecular characterization of *Withania* accessions and establishing genetic relatedness among genetic stock evaluated. High information content of the markers enables differentiation and discrimination of *Withania* accessions. The results are in accordance with those of Ben Hmida-Ben Salem (2000) who found that SSR markers seemed to be effective to discriminate local barleys defined as accessions or populations geographically based. Yu et al. (2012) examined the genetic diversity of 80 inbred waxy maize lines using 22 SSR molecular markers that could be used to achieve heterosis in waxy maize and found the values genetic distance from (0.233 to 0.505). An SSR marker used for diversity studies has recorded genetic distance range of 16-53% among 14 genotypes of maize using four SSR markers (Bibi et al., 2010). These results are in agreement with results of the present study.

SSR data has separated the accessions into two major genetic similarity groups. These findings have direct correlation with chemical profiling data. This correlation indicate that the independent sets of data reflect the same pattern of genetic diversity and validate the use of these methods for diversity estimation and also in grouping of accessions. The results also indicate that the combination of morphological and molecular markers may be useful in studying genetic diversity as reported by Cortese et al. (2010). Our results are also in conformity with Roldan-Ruiz et al. (2001) reported that when varieties with shared genepools were examined using molecular markers, extremely high similarity measures were produced and were also linked to morphological similarities. SSRs are relatively quick and easy to use, but refractory to many environmental influences. However, they can be generated in large numbers and can complement traditional character that may be limited in availability. SSRs also provide a valuable new resource for phylogenetic studies. The results obtained in this study may assist *Withania* cultivation and in *Withania* breeding programmes.

The combination of microsatellite marker profiles, chemical and the morphological characters serve as reliable tools for detailed description of accessions. Identification of superior genotypes by assessing genetic diversity is an important prerequisite for a successful crop improvement program.

Acknowledgement

PKK is thankful to Shri Mata Vaishno Devi University for providing research facility and opportunity to pursue the Ph.D. programme.

References

- Akkaya MS, Bhagwatt AA, Cregan PB. 1992. Length polymorphism of simple sequence repeat DNA in soybean. *Genetics*, 132: 1131-1139.
- Atta-ur-Rahman, Yousaf M, Gul W, Qureshi S, Choudhary MI, Voelter W, Hoff A, Jens F, Naz A. 1998. Five new withanolides from *Withania coagulans*. *Heterocycles*, 48: 1801-1811.
- Ayres NM, McClung AM, Larkin PD, Bligh HFJ, Jones CA, Park, WD. 1997. Microsatellite and a single nucleotide polymorphism differentiate apparent amylose classes in an extended pedigree of US rice germplasm. *Theor. Appl. Genet.*, 94: 773-781.
- Bandyopadhyay M, Jha S. 2003. *Withania somnifera* (Ashwagandha): A review. *Journal of Tropical Medicinal Plants*, 4: 273-284.
- Ben Hmida-Ben Salem W. 2000. Contribution des marqueurs microsatellites et RAPD pour l'études du polymorphisme chez l'orge. Diplômes d'étude approfondies de génétique et iologie moléculaires, Faculté. Sciences, Tunis, Tunisie, 80 p.
- Benor S, Zhang MY, Wang ZF, Zhang HS. 2008. Assessment of genetic variation in tomato (*Solanum lycopersicum* L.) inbred lines using SSR molecular markers. *Journal of Genetics and Genomics*, 35: 373-379.
- Bhattacharya SK, Muruganandam AV. 2003. Adaptogenic activity of *Withania somnifera*: an experimental study using a rat model of chronic stress. *Pharmacol. Biochem. Behav.*, 75: 547-555.
- Bhattacharya SK, Bhattacharya D, Sairam K, Ghosal S. 2002. Effect of *Withania somnifera* glycowithanolides on a rat model of tardive dyskinesia. *Phytomedicine*, 9 (2): 167-170.
- Bibi S, Ahmad H, Ghafoor S, Begum K, Afridi SG, Khan IA. 2010. Genetic analysis of some commonly grown genotypes of maize in Pakistan using maize specific simple sequence repeat (SSR) markers. *Current Res. J. Bio. Sci.*, 2(4): 259-261.
- Bone K. 1996. Clinical applications of Ayurvedic and Chinese herbs. Monographs for the western herbal practitioner. Australia: Phytotherapy press, 137-141.
- Chandra A. 2011. Use of EST database markers from *Medicago truncatula* in the transferability to other forage legumes. *J. Environ. Biol.*, 32: 347-354.
- Cortese LM, Honig J, Miller C, Bonos SA. 2010. Genetic diversity of twelve switchgrass populations using molecular and morphological markers. *Bioenergy Research*, 3: 262-271.
- Datta S, Kaashyap M, Kumar S. 2010a. Amplification of chickpea-specific SSR primers in *Cajanus* species and their validity in diversity analysis. *Plant Breed.*, 129: 334-340.
- Datta S, Mahfooz S, Singh P, Choudhary AK, Singh F, Kumar S. 2010b. Cross-genera amplification of informative microsatellite markers from common bean and lentil for the assessment of genetic diversity in Pigeon pea. *Physiol. Mol. Biol. Plants*, 16: 123-134.

RESEARCH ARTICLE

- Datta S, Tiwari S, Kaashyap M, Gupta PP, Choudhury PR, Kumari J, Kumar S. 2011. Genetic similarity analysis in lentil using cross-genera legume sequence tagged microsatellite site markers. *Crop Sci.*, 51: 2412-2422.
- Dhuley JN. 2000. Adaptogenic and cardioprotective action of ashwagandha in rats and frogs. *J. Ethnopharmacol.*, 70(1): 57-63.
- Diwan N, Cregan PB. 1997. Automated sizing of fluorescent-labeled simple sequence repeat markers to assay genetic variation in soybean. *Theor. Appl. Genet.*, 95: 723-733.
- Doyle JJ, Doyle JL. 1990. Isolation of plant DNA from fresh tissue. *Focus*, 12: 13-15.
- Eujay I, Sorrells M, Baum M, Woltersand P, Powell W. 2001. Assessment of genotypic variation among cultivated durum wheat based on EST-SSRs and genomic SSRs. *Euphytica*, 119: 39-43.
- Garcia-Vallve S, Palau J, Romeu A. 1999. Horizontal gene transfer in glycosyl hydrolases inferred from codon usage in *Escherichia coli* and *Bacillus subtilis*. *Mol. Biol. Evol.*, 16: 1125-1134.
- Grierson DS, Afolayan AJ. 1999. Antibacterial activity of some indigenous plants used for the treatment of wounds in the Eastern Cape, South Africa. *J. Ethnopharmacol.*, 66: 103-106.
- Grover A, Ramesh B, Sharma PC. 2009. Development of microsatellite markers in potato and their transferability in some members of Solanaceae. *Physiol. Mol. Biol. Plants*, 15: 343-358.
- Hamza S, Hamida WB, Rebai A, Harrabi M. 2004. SSR based genetic diversity among Tunisian Barley and relationship with morphological traits. *Euphytica*, 135: 107-118.
- Jaccard P. 1901. Étude comparative de la distribution florale dans une portion des Alpes et des Jura. *Bull. Soc. Vandoise Sci. Nat.*, 37: 547-579.
- Jayaprakasam B, Zhang Y, Seeram NP, Nair MG. 2003. Growth inhibition of human tumor cell lines by withanolides from *Withania somnifera* leaves. *Life Sci.*, 74: 125-132.
- Khajuria RK, Suri KA, Gupta RK, Satti NK, Amina M, Suri OP, Qazi GN. 2004. Separation, identification and quantification of selected withanolides in plant extracts of *Withania somnifera* by HPLC-UV(DAD) – Positive ion electrospray ionization-mass spectrometry. *J. Sep. Sci.*, 27: 541-546.
- Mary NK, Babu BH, Padikkala J. 2003. Antiatherogenic effect of Caps HT2, a herbal Ayurvedic medicine formulation. *Phytomedicine*, 10: 474-482.
- Matsuda H, Murakami T, Kishi A, Yoshikawa M. 2001. Structures of withanosides I, II, III, IV, V, VI and VII new withanolide glycosides from the roots of Indian *Withania somnifera* Dunal and inhibitory activity for tachyphylaxis to clonidine in isolated guinea-pig ileum. *Bioorg. Med. Chem.*, 9: 1499-1507.
- Maughan PJ, Saghai-Maroo MA, Buss GR. 1995. Microsatellite and amplified sequence length polymorphism in cultivated and wild soybean. *Genome*, 38: 715-723.
- Mishra LC, Singh BB, Dagenais S. 2000. Scientific basis for therapeutic use of *Withania somnifera* (ashwagandha): a review. *Altern. Med. Rev.*, 5: 334-346.
- Mohammadi SA, Prasanna BM. 2003. Analysis of genetic diversity in crop plants-salient statistical tools and considerations. *Crop Sci.*, 43: 1235-1248.
- Peakall R, Gilmore S, Keys W, Morgante M, Rafalski A. 1998. Cross-species amplification of Soybean (*Glycine max*) simple sequence repeats (SSRs) within the genus and other legume genera: implication for the transferability of SSRs in plants. *Mol. Biol. Evol.*, 15: 1275-1287.
- Pillen K, Binder A, Kreuzkam B, Ramsay L, Waugh R, Förster J, Leon J. 2000. Mapping new EMBL-derived barley SSRs and their use in differentiating German barley cultivars. *Theor. Appl. Genet.*, 101: 652-660.
- Rafalski JA, Vogel JM, Morgante M, Powell W, Andre C, Tingey SV. 1996. Generating and using DNA markers in plants. - In: Birren B. & Lai E. (eds), *Nonmammalian genome analysis-a practical guide*, Academic Press Inc, San Diego, p. 75-135.
- Roldan-Ruiz I, van Eeuwijk FA, Gilliland TJ, Dubreuil P, Dillmann C, Lallemand J, De Loose M, Baril CP. 2001. A comparative study of molecular and morphological methods of describing relationships between perennial rye grass (*Lolium perenne* L.) varieties. *Theor. Appl. Genet.*, 103: 1138-1150.
- Rongwen J, Akkaya MS, Bhagwat AA, Levi U, Cregan PB. 1995. The use of microsatellite DNA markers for soybean genotypes identification. *Theor. Appl. Genet.*, 90: 43-48.
- Russell J, Booth A, Fuller J, Harrower B, Hedley P, Machray G, Powell W. 2004. A comparison of sequence-based polymorphism and haplotype content in transcribed and anonymous regions of the barley. *Genome*, 47: 389-398.
- Saghai-Maroo MA, Biyashev RM, Yang GP, Zhang Q, Allard RW. 1994. Extraordinarily polymorphic microsatellite DNA in barley: species diversity, chromosomal locations, and population dynamics. *Proc. Natl. Acad. Sci. USA*, 91: 5466-70.
- Saker M, Nagchtigall M, Kuehne TA. 2005. Comparative assessment of DNA fingerprinting by RAPD, SSR and AFLP in genetic analysis of some barley genotypes. *Egypt. J. Genet. Cytol.*, 34: 81-97.
- Sharma R, Mahla HR, Mohapatra T, Bhargava SC, Sharma MM. 2003. Isolating plant genomic DNA without liquid nitrogen. *Plant Mol. Biol. Rep.*, 21: 43-50.
- Singh A, Naidu PS, Gupta S, Kulkarni SK. 2002. Effect of natural and synthetic antioxidants in a mouse model of chronic fatigue syndrome. *J. Med. Food.*, 5(4): 211-220.
- Smith DN, Devey ME. 1994. Occurrence and inheritance of microsatellites in *Pinus radiata*. *Genome*, 37: 977-983.
- Somta P, Seehalak W, Srinives P. 2009. Development, characterization and cross-species amplification of Mungbean (*Vigna radiata*) genic microsatellite markers. *Conserv. Genet.*, 10: 1939-1943.
- Thompson JA, Nelson RL, Vodkin LO. 1998. Identification of diverse Soybean germplasm using RAPD markers. *Crop Sci.*, 38: 1348-1355.
- Vosman B, Arens P, Rus-Kortekaas W, Smulders MJM. 1992. Identification of highly polymorphic DNA regions in tomato. *Theor. Appl. Genet.*, 85: 239-244.
- Yu RH, Wang YL, Sun Y, Liu B. 2012. Analysis of genetic distance by SSR in waxy maize. *Genetics and Molecular Research*, 11(1): 254-260.